

A cytochrome P450 regulates a domestication trait in cultivated tomato

Manohar Chakrabarti^{a,1}, Na Zhang^{a,1,2}, Christopher Sauvage^b, Stéphane Muñoz^c, Jose Blanca^d, Joaquin Cañizares^d, Maria Jose Diez^d, Rhiannon Schneider^a, Michael Mazourek^e, Jammi McClead^a, Mathilde Causse^b, and Esther van der Knaap^{a,3}

^aDepartment of Horticulture and Crop Science, The Ohio State University/Ohio Agricultural Research and Development Center, Wooster, OH 44691; ^bInstitut National de la Recherche Agronomique (INRA), Unité de Génétique et Amélioration des Fruits et Légumes, Domaine Saint-Maurice, F-84143 Montfavet Cedex, France; ^cCentre National de la Recherche Scientifique-INRA, Unité Mixte de Recherche 2594/441, 31326 Castanet-Tolosan Cedex, France; ^dInstitute for the Conservation and Improvement of Agricultural Biodiversity, Polytechnic University of Valencia, Valencia 46022, Spain; and ^eDepartment of Plant Breeding and Genetics, Cornell University, Ithaca, NY 14853

Edited by James J. Giovannoni, US Department of Agriculture-Agricultural Research Service Robert W. Holley Center and Boyce Thompson Institute for Plant Research, Ithaca, NY, and accepted by the Editorial Board August 23, 2013 (received for review April 22, 2013)

Domestication of crop plants had effects on human lifestyle and agriculture. However, little is known about the underlying molecular mechanisms accompanying the changes in fruit appearance as a consequence of selection by early farmers. We report the fine mapping and cloning of a tomato (*Solanum lycopersicum*) fruit mass gene encoding the ortholog of KLUH, *SIKLUH*, a P450 enzyme of the CYP78A subfamily. The increase in fruit mass is predominantly the result of enlarged pericarp and septum tissues caused by increased cell number in the large fruited lines. *SIKLUH* also modulates plant architecture by regulating number and length of the side shoots, and ripening time, and these effects are particularly strong in plants that transgenically down-regulate *SIKLUH* expression carrying fruits of a dramatically reduced mass. Association mapping followed by segregation analyses revealed that a single nucleotide polymorphism in the promoter of the gene is highly associated with fruit mass. This single polymorphism may potentially underlie a regulatory mutation resulting in increased *SIKLUH* expression concomitant with increased fruit mass. Our findings suggest that the allele giving rise to large fruit arose in the early domesticates of tomato and becoming progressively more abundant upon further selections. We also detected association of fruit weight with *CaKLUH* in chile pepper (*Capsicum annuum*) suggesting that selection of the orthologous gene may have occurred independently in a separate domestication event. Altogether, our findings shed light on the molecular basis of fruit mass, a key domestication trait in tomato and other fruit and vegetable crops.

Plant domestication and artificial selection led to improved agricultural production resulting from dramatic increases in fruit and seed weight (1). At the start of the Neolithic era ~10,000 y ago, domestication of animals and plants accompanied the change in lifestyle from hunter-gatherer to a farming routine (2, 3). Driven by the selection of alleles from wild relatives and those that arose after the initial domestication events, characters associated with the domestication syndrome such as larger fruit and seed are typically differentiating wild from cultivated forms (4). Genome-wide genetic diversity analyses support the notion that the initial domestication of tomato was situated in Northern Peru and Ecuador (5). Selections from the red-fruited wild relative *Solanum pimpinellifolium* L. evolved into the semidomesticated *Solanum lycopersicum* L. var *cerasiforme* (hereafter referred to as *S. l. cerasiforme*) bearing fruit of small to medium weight. *S. l. cerasiforme* was further domesticated in Mexico giving rise to the large fruited tomato *Solanum lycopersicum* var *lycopersicum* (hereafter referred to as *S. l. lycopersicum*), which to date is cultivated throughout the world (5).

Tomato is an agriculturally important vegetable crop and is used as model for fruit development including ripening and morphological studies (6–8). Genetic studies have identified several quantitative trait loci (QTLs) associated with fruit mass in tomato, of which six loci [*fruit weight1.1* (*fw1.1*), *fw2.2*, *fw2.3*,

fw3.1/fw3.2, *fw4.1*, and *fw9.1*] are postulated to be major QTLs (9). The only cloned fruit mass gene from vegetable or fruit crops is *FW2.2/Cell Number Regulator* (*CNR*) (10). The gene encodes a negative regulator of cell division and controls tomato fruit mass as well as organ size in other species, e.g., maize (10, 11) and nitrogen-fixing nodule number (12). *fw3.2* is the second major tomato fruit mass QTL, which explained 19% of the phenotypic variance in a F₂ population derived from a cross between cultivated tomato accession “Yellow Stuffer” and wild tomato accession “LA1589” (13, 14). Despite the importance of fruit mass in the evolution of fruit and vegetable plants, and the numerous genetic loci that underlie the trait (7, 9, 15–19), cloning of domestication genes of fruit and vegetable crops has lagged behind that of the cereal crops. Therefore, insights into the molecular mechanisms that led to the transition of the fruit from small to large remain largely unknown.

Our current study focuses on fine mapping of a tomato fruit mass locus *fw3.2* and cloning of the underlying gene. Association mapping, segregation analysis, and transgenic studies led us to identify the putative molecular basis of fruit weight at this locus and a likely regulatory SNP in the promoter of the gene that is highly associated with fruit mass. Phenotypic

Significance

This study reports the cloning of a tomato gene, *SIKLUH*, that controls fruit mass by increased cell layers and delayed fruit ripening. In addition, we identified a potential regulatory SNP in the promoter of *SIKLUH* that is significantly associated with the fruit mass. Altogether, our study encompasses several genetic analyses, as well as association mapping, plant transformation experiments, and phenotypic evaluations to offer insights into the molecular basis of the regulation of tomato fruit mass, a critical trait in the domestication of fruit and vegetable crops.

Author contributions: E.v.d.K. designed research; M. Chakrabarti, N.Z., C.S., S.M., J.B., J.C., R.S., M.M., J.M., and E.v.d.K. performed research; M.J.D., M.M., M. Causse, and E.v.d.K. contributed new reagents/analytic tools; M. Chakrabarti, N.Z., C.S., S.M., J.B., J.C., and E.v.d.K. analyzed data; and M. Chakrabarti, N.Z., and E.v.d.K. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission. J.J.G. is a guest editor invited by the Editorial Board.

Data deposition: The sequences reported in this paper have been deposited in the National Center for Biotechnology Information Sequence Read Archive (SRA submission nos. SRA068200 and SRA091611) and in the Genomic Survey Sequences (GSS accession nos. KG700052–KG701213).

¹M.C. and N.Z. contributed equally to this work.

²Present address: Taiyuan Research Center/DuPont Pioneer, Jinzhong, Shanxi 030622, China.

³To whom correspondence should be addressed. E-mail: vanderknaap.1@osu.edu.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1307313110/-DCSupplemental.

Table 1. Effect of *fw3.2* on fruit attributes and ripening

Measurements	<i>fw3.2</i> (ys)	<i>fw3.2</i> (wt)	P value
Perimeter, cm			
Ovary	0.61 ± 0.05	0.61 ± 0.06	0.75900
Mature fruit	19.56 ± 0.51	16.68 ± 0.30	9.6E-08
Area of mature fruit parts, cm ²			
Columella and Placenta	3.05 ± 0.35	2.65 ± 0.19	0.01475
Septum	2.62 ± 0.34	1.40 ± 0.23	1.5E-05
Pericarp	11.48 ± 0.47	8.36 ± 0.36	1.6E-07
Pericarp cell size, μm			
Ovary	7.23 ± 0.62	7.64 ± 0.93	0.32222
Mature fruit	616.67 ± 68.12	624.63 ± 54.65	0.74632
Pericarp cell number			
Ovary	9.04 ± 0.52	9.26 ± 0.49	0.40223
Mature fruit	19.20 ± 1.75	17.07 ± 0.93	0.00458
Fruit ripening, days			
Anthesis to orange	48.87 ± 1.06	43.14 ± 0.38	8.1E-08
Anthesis to red	53.44 ± 0.97	47.27 ± 1.11	3.2E-08

Data were taken from 6 to 10 plants per genotype and are given as mean ± SD.

evaluations demonstrated the cellular basis of increased fruit mass and pleiotropic effects associated with the locus. We also investigated the likely origin of the derived allele and the potential role of this gene in the regulation of fruit mass in other crop species.

Results

***fw3.2* Regulates Fruit Mass, Delays Ripening, and Modulates Plant Architecture.** To elucidate whether fruit mass differences were determined during flower or fruit development, we evaluated the ovary size at anthesis. Evaluations of the nearly isogenic lines (NILs) that differ for the allele at *fw3.2* showed that the ovary perimeter at anthesis was similar, whereas the mature fruit perimeter was significantly different. Further analyses revealed that in particular the pericarp and septum areas were significantly larger in the large fruited NIL *fw3.2*(ys) (Table 1 and Fig. S1 A and B). The pericarp of the mature fruit carrying the large fruit allele *fw3.2*(ys) showed an increase in cell number, whereas cell size remained the same (Table 1 and Fig. S1 C and D). The time from anthesis to ripe fruit was prolonged in the *fw3.2*(ys) NIL (Table 1). In addition to fruit, *fw3.2*(ys) increased seed mass, although seed number was not significantly different (Table S1). Number of fruits were significantly higher in the lines carrying the small fruited *fw3.2*(wt) allele, whereas yield per plant was the same (Table 2). Additionally, *fw3.2*(wt) NIL contained more inflorescences and side shoots, including increased side shoot length (Table 2 and Table S1). The number of flowers per inflorescence remained the same. Retaining only 12 fruits per plant led to increased fruit weight in both NILs (Table S1) demonstrating that the increase in fruit mass controlled by *fw3.2* was not due to changes in source–sink relationships. Thus, the results demonstrate that increases in fruit mass coincide with a reduction of number of fruit per plant caused by a reduction in side shoot number and length yielding fewer inflorescences.

Fine Mapping Delimited the *fw3.2* Locus to a 24.4-kb Region. Previously, the *fw3.2* locus was fine mapped to a 51.4-kb region comprised of seven candidate genes (14). An additional recombinant screen delimited the locus to a 24.4-kb region comprised of three candidate proteins: a cytochrome P450 (*ORF6*) belonging to CYP78A subfamily, an ABC transporter (*ORF7*) distantly related to PGP transporters, and a Kelch domain-containing protein of unknown function (*ORF8*). The fine-mapping experiment excluded most of the *ORF8* gene to only include the promoter, first exon, and first intron until the NDF9 marker (Fig. 1 and Table S2). Gene

action analysis revealed the additive nature of the *fw3.2* alleles, suggesting that neither allele is a null (Table S2).

Association Mapping and Genetic Segregation Analysis Identified a Potential Regulatory SNP in the Promoter of *ORF6*. To further investigate which of the candidate genes may underlie the *fw3.2* locus, we sought to identify the underlying nucleotide polymorphisms that were correlated with larger fruit. We conducted an association mapping study using a core collection of tomato genotypes demonstrated to have broad genetic diversity within red-fruited tomatoes and comprised of domesticated *S. l. lycopersicum*, the wild relative *S. pimpinellifolium* and semidomesticated *S. l. cerasiforme* accessions (20). We sequenced ~7 kb of the *fw3.2* locus and genotyped 115 markers inside and outside the *fw3.2* region spanning 48.5 kb (Fig. 1 and Dataset S1). Accounting for kinship and population structure, we identified six SNPs that were significantly associated with fruit mass (Fig. 1 and Table S3). With the exception of one SNP, all were located in the upstream region of *ORF6* and *ORF7*. The two most significant SNPs were found in or near *ORF6*, 512 bp upstream of the start of transcription (M9) and 72 bp from the stop codon in the 3'UTR (M42) (Fig. 1). Successful association mapping typically requires rapid linkage disequilibrium (LD) decay. However, low rates of LD decay were found around the *fw3.2* locus, suggesting few natural recombination events in this region that would potentially compromise the interpretation of the association mapping results (21). Therefore, to determine the validity of the association of these SNPs to fruit mass, we performed intraspecific segregation analyses using populations derived from parents that showed few polymorphisms including M9 and three to five additional SNPs (Dataset S1 and Table S4). Genetic analyses using the F₂ and BC₁F₂ populations showed segregation for fruit mass with the allele of M9 in all families in either one or both populations (Table S4). Importantly, family 12S74 segregated for M9 but not for M42 and showed that fruit mass segregated with the allele of M9 (Dataset S1 and Table S4). This finding demonstrated that the M9 SNP was more critical in regulating fruit mass than M42, lending support for the notion that the *ORF6* promoter SNP may regulate fruit mass differences at *fw3.2*.

Table 2. Effect of *fw3.2* on yield parameters

Yield parameters	<i>fw3.2</i> (ys)	<i>fw3.2</i> (wt)	P value
Plant height, cm			
At 35 das	6.00 ± 1.94	5.50 ± 1.65	0.55705
At 56 das	59.00 ± 10.25	54.20 ± 8.79	0.12792
No. of nodes			
At 35 das	7.10 ± 1.20	7.90 ± 0.74	0.15266
At 56 das	17.90 ± 1.79	19.40 ± 1.07	0.07138
No. of side shoots			
At 56 das	7.11 ± 1.36	10.80 ± 1.23	1.3E-05
At 70 das	9.67 ± 1.66	12.90 ± 0.99	0.00022
Total side shoot length, cm			
At 56 das	116.25 ± 42.47	177.85 ± 41.93	0.01761
At 70 das	283.60 ± 65.24	405.95 ± 50.14	0.00346
No. of fruits			
Red fruits	175.30 ± 42.28	257.90 ± 77.70	0.01582
Green fruits	163.90 ± 40.06	209.40 ± 57.46	0.01036
Total fruits	339.20 ± 71.19	467.30 ± 126.60	0.00600
Total fruit weight per plant, kg			
Red fruits	10.17 ± 3.10	10.44 ± 3.87	0.85921
Green fruits	5.66 ± 1.88	5.00 ± 1.69	0.20173
Total yield	15.83 ± 4.56	15.44 ± 5.37	0.83032

Data were collected from 10 plants per genotype and are given as mean ± SD das, days after sowing.




Fig. 1. Genome structure of the *fw3.2* locus and association mapping of the polymorphisms. UTRs, exons, introns, and intergenic regions are represented with red boxes, black boxes, black lines, and green lines, respectively. The direction of transcription is denoted with red arrows. Blue vertical lines represent polymorphisms not associated with fruit mass; red vertical lines represent polymorphisms significantly associated with fruit mass. Two highly significant SNPs are represented with an asterisk. Red horizontal lines depict the sequenced regions. Two markers, P450-2 and NDF9, which delimited the *fw3.2* locus to 24.4-kb region, are shown with green vertical lines.

Down-Regulation of *ORF6* Leads to Reduced Fruit Mass. To further investigate the role of the candidate genes in regulating fruit mass, we down-regulated transcript accumulation of *ORF6* and *ORF7*. Because expression of *ORF6* and *ORF7* was lower in the lines that carried small fruit (Fig. 2A), the constructs were transformed into the *fw3.2*(*ys*) NILs typically yielding large fruit. Down-regulation of *ORF6* transcript levels dramatically reduced fruit and seed mass (Fig. 2B–E and Table S5). Similar to the NILs, down-regulation of *ORF6* led to a higher number of side shoots and faster ripening (Table S5). The transgenic lines showed additional defects compared with the NILs (e.g., reduced plant height, smaller leaves and leaflets, and severely reduced seed number), indicating that extensive down-regulation of *ORF6* led to phenotypes affecting the entire plant (Table S5). The RNAi-2 lines targeting the coding region led to less severe phenotypes, including slightly larger fruit with viable seeds. Even though overall seed number was significantly reduced, fruit with 50–100 seeds showed reduced fruit weight compared with the control with the same number of seed (Fig. S1H). Thus, when accounting for seed number, fruit mass is still reduced in lines that down-regulate *ORF6*. Contrary to *ORF6*, down-regulation of *ORF7* using four amiRNA constructs did not show a correlation between reduction in fruit mass and transcript level (see Table S7 and Fig. S1I), and the plants were indistinguishable from the *fw3.2*(*ys*) NIL. Thus, it is unlikely that *ORF7* underlies the *fw3.2* locus. In light of these findings, we assumed that *ORF6* underlies the *fw3.2* locus and renamed it *SIKLUH* after one of the founding members of this subfamily of P540s (22). Expression of *SIKLUH* was the highest in vegetative meristems and in the young flower buds (Fig. 2F). Compared with *SIKLUH*, the expression of *SIFW2.2/CNR* was much lower in most tissues examined and differed in tissue specificity. Also, expression of *SIKLUH* was very high in developing seed and not in the growing pericarp, contrary to *SIFW2.2/CNR* (Fig. 2G).

Molecular Diversity and Phylogenetic Analysis of *SIKLUH*. To examine the evolutionary history of the *fw3.2* locus, DNA sequence variation of the segments spanning part of the coding region (fragment A) and promoter (fragment B) of *SIKLUH* were investigated. The *SIKLUH* region showed reduced nucleotide diversity (π) in *S. lycopersicum* (0 and 0.000195 for fragment A and B, respectively) compared with *S. pimpinellifolium* (0.002874 and 0.004962) and *S. l. cerasiforme* (0.003203 and 0.004364) (Fig. 3A). The ratio of nucleotide diversity of *S. l. lycopersicum* to *S. pimpinellifolium* at *SIKLUH* was significantly lower compared with the average diversity for chromosome 3 [percentage of $\pi_{S. lycopersicum}/\pi_{S. pimpinellifolium}$ for fragment A, B, and chromosome 3 are 0%, 3.9%, and 56.9%, respectively]. Tajima's *D* analysis of all accessions combined showed significant values for parts of the two sequenced fragments, ranging from -2.0851 to -1.5187 and from -2.1922 to -1.9117 for fragment A and B, respectively, compared with the average Tajima's *D* for chromosome 3 (-1.3247) (Fig. 3A). The findings of the molecular diversity analysis were also supported by the results from the phylogenetic analysis using the *SIKLUH* gene sequences. One cluster consisted of all *S. pimpinellifolium* and several *S. l. cerasiforme* accessions and

displayed relatively high sequence diversity. The other cluster is comprised of the *S. l. lycopersicum* and the remaining *S. l. cerasiforme* accessions and showed limited genetic diversity (Fig. 3B). The derived allele designated by the M9 SNP was only found in a single subclade with some *S. l. cerasiforme* and all of the cultivated tomato accessions, suggesting the mutation arose in the *S. l. cerasiforme* background.

Possible Origin of the M9 Mutation and Identification of Fruit Mass QTL Overlapping with *fw3.2* in Chile Pepper. To further explore where the derived allele arose, we genotyped the M9 SNP in a tomato collection that covers the proposed trajectory of tomato domestication (5). The M9 SNP associated with increased fruit mass was already present at $\sim 30\%$ frequency in the Ecuadorian and Northern Peruvian *S. l. cerasiforme* that were closely related to the wild progenitor *S. pimpinellifolium*. The allele frequency increased slightly in the Mesoamerican *S. l. lycopersicum* accessions, and increased further in the *S. l. lycopersicum* landraces from Europe and elsewhere to become practically fixed in modern accessions (Fig. 3C and Table S6). This finding suggested that even though the derived allele arose early, selections for larger fruit weight may not have been as critical or this allele was not selected early and only became relevant in recent times.

Based on coinciding QTLs, it has been proposed that orthologous genes may have been selected independently in different crops




Fig. 2. Transcript accumulation and effect of *SIKLUH* knock down on fruit and seed mass. (A) Expression of *ORF6* (*SIKLUH*) and *ORF7* in *fw3.2*(*ys*) and *fw3.2*(*wt*) NILs. (B) Regions of *SIKLUH* targeted by two RNAi constructs. (C and D) Effect of *SIKLUH* knock down on fruit and seed mass, respectively. (E) Transcript accumulation of *SIKLUH* in *fw3.2*(*ys*) and *fw3.2*(*wt*) NILs and *SIKLUH* knock down lines. (F) Average transcript accumulation of *SIKLUH* and *SIFW2.2/CNR* in different tissues of *S. pimpinellifolium* LA1589. (G) Average transcript accumulation of *SIKLUH* and *SIFW2.2/CNR* in *S. l. lycopersicum* pericarp and developing seeds at different fruit development stages. dpa, days postanthesis; P, pericarp; S, seed; Veg. meristem, vegetative meristem.




Fig. 3. Molecular diversity and phylogenetic analysis of *SIKLUH*, and origin of M9 mutation. (A) Nucleotide diversity (π) and Tajima's D were calculated in fragment A and B spanning part of the coding region and promoter of *SIKLUH* (Fig. 1). The Tajima's D is calculated from the *SIKLUH* sequences of *S. pimpinellifolium* (*S. p.*), *S. l. cerasiforme* (*S. l. c.*), and *S. lycopersicum lycopersicum* (*S. l. l.*) combined. (B) Phylogenetic analysis of *SIKLUH*. *S. l. l.*, *S. l. c.*, *S. p.*, and *Solanum pennellii* are depicted in red, black, green, and blue, respectively. Large (>60 g), medium (>10 g and <60 g), and small (<10 g) fruit-bearing lines are represented as large, medium, and no dots after their name, respectively. Accessions carrying the M9 mutation are depicted with an asterisk (*). (C) Frequency of derived and ancestral M9 SNP allele in tomato subpopulations. Imp, improved accessions; Ecu, Ecuador; Mes, Mesoamerica; N.Per, Northern Peru; Per, Peru.

(7, 23). To evaluate whether chile pepper (*Capsicum annuum*) may harbor a fruit mass QTL at a region that is syntenic to tomato *fw3.2*, we determined whether the pepper ortholog *CaKLUH* was associated with fruit mass in this species. A population of 106 F9 recombinant inbred lines (RILs) derived from a cross between a large-fruited pepper and a small-fruited wild relative *Capsicum frutescens* showed fruit mass segregated significantly with the large fruited allele of *KLUH* from the *C. annuum* parent [dry fruit weight (mean \pm SD, g) for groups carrying the *C. frutescens* and the *C. annuum* allele of *fw3.2* are 3.21 ± 0.36 and 4.99 ± 0.34 , respectively; P value < 0.0005]. These findings demonstrate that *fw3.2* QTL is found in both tomato and chile pepper, and that *KLUH* may regulate fruit mass in two independently domesticated species.

Discussion

***SIKLUH* Is Most Likely Underlying *fw3.2*, and a Critical SNP in Its Promoter Is Proposed to Regulate Fruit Mass.** *ORF6* encodes a protein with the highest similarity to the *Arabidopsis* *KLUH*/CYP78A5 protein controlling vegetative and reproductive organ size, and also seed size (22, 24). Other members of this subfamily of cytochrome P450 also control organ and plant size, e.g., CYP78A9, CYP78A6

in *Arabidopsis*, and CYP78A11 in rice (25–27). Two additional members of this subfamily, CYP78A27 and CYP78A28, control colony size in moss (28). This highly conserved function of CYP78A subfamily in regulating organ size led us to postulate that *ORF6* was the most likely candidate gene for the *fw3.2* QTL.

The first experiment to test this hypothesis was to identify SNPs that were significantly associated with fruit mass. Our expectations were that these SNPs would be found near the most likely candidate gene. The most significant SNPs identified from the association mapping were the M9 SNP located 512 bp upstream of the transcription start site and the M42 SNP located in the 3' UTR at 72 bp downstream from the stop codon of *ORF6*. However, LD decay rate was low (21), which was likely due to the lack of outcrossing and recombination events during domestication of tomato. However, when the M9 polymorphism arose, its effect on fruit mass could have been of functional significance. This pattern of few SNPs in large LD blocks has also been found in human (29–32) and may be common in plants as well. The low LD yet significant association of the SNP with fruit mass led us to conduct segregation analyses; these analyses excluded the importance of M42, leaving M9 as the most significantly associated SNP of potential functional relevance (i.e., the most likely to regulate fruit mass). Interestingly, the M9 SNP was located within the third repeat of four 30-bp tandem repeats, with each repeat sharing 83–100% homology with its consensus sequence. The tandem repeat was located 440 bp to 559 bp upstream from the *SIKLUH* transcription start site. Moreover, M9 mutation is located within a putative *cis* element, related to the organ-specific element (OSE) found in nodulin and leghemoglobin genes in soybean and *Vicia faba* (33–35). In the future, it will be interesting to assess the role of this putative *cis*-element in gene expression and to explore whether the tandem repeat has any bearing on the modulation of *SIKLUH* expression. Even though the M9 SNP appears most promising to regulate fruit mass, we cannot exclude the roles of three other SNPs (M60, M74, and M82) in the regulation of fruit mass, and additional SNPs that may be found in the remaining regions that were not sequenced cannot be excluded either.

The second experiment was to transgenically down-regulate the expression of the candidate genes, *ORF6* and *ORF7*. Our expectation was that expression levels positively correlate with fruit mass since the *fw3.2(wt)* NIL exhibit smaller fruit and lower transcript levels. Reduced expression of *ORF6*, and not of *ORF7*, had pronounced effects on fruit mass, seed mass, side shoot number, and fruit ripening, which were traits that were also significantly altered in the NILs. The severity of these phenotypes followed a gradient from *fw3.2(ys)*, *fw3.2(wt)* to the RNAi line. Thus, fine mapping, association mapping, and functional segregation analyses of a highly significant SNP in the promoter of *ORF6* combined with the plant transformation results and phenotypic evaluations strongly implied that *ORF6* underlies the *fw3.2* locus. The expression of *ORF6*, which we renamed *SIKLUH*, was particularly high in the developing seeds. However, the effect of increased fruit mass was found in the pericarp of the maternal tissues. Thus, it is plausible that a seed-derived signal might play a role in regulation of fruit mass by *SIKLUH*. A similar hypothesis has been proposed to explain the action of *Arabidopsis* *KLUH* (22).

The Role of *SIKLUH* in Regulating Fruit Mass, Ripening, and Plant Architecture. Fruit development starts with the fertilization of the ovules in the ovary at the time of anthesis (flower opening) leading to the initiation of seed development. Immediately following fertilization, the fruit undergoes a short period of cell division, which is followed by a longer period of cell expansion until the final dimensions are reached (36–39). The increase in fruit mass was due to growth processes taking place after fertilization, particularly of the pericarp and septum tissues. These findings showed that the extra cell divisions led to enlarged fruit and a concomitant delay in ripening. Therefore, the delay in ripening was likely the result of an extension of the cell division stage resulting from increased expression. *SIFW2.2/CNR* also

controls fruit mass by extending the cell division period (40). Thus, the increases in fruit mass resulting from domestication may operate on similar processes during fruit development yet through different mechanisms based on putative gene function. The lengths of other plant organs were not affected in the NILs, suggesting a specific role of the natural alleles of *SIKLUH* in the regulation of fruit and seed mass. Our results also indicated that plant architecture was affected, implying a pleiotropic effect of *SIKLUH* on plant growth and development.

Diversity of *SIKLUH* and Origin of M9 Mutation. Our findings indicate reduced nucleotide diversity in *S. l. lycopersicum* in the *SIKLUH* region compared with *S. pimpinellifolium*, and an overall reduced diversity compared with the entire chromosome. Parts of two sequenced regions around *SIKLUH* showed significant Tajima's *D* values, which may suggest an excess of low frequency SNPs in certain regions of the gene in tomato, which is potentially the result of a population expansion after a bottleneck or a selective sweep. Taken together, these results support a selective pressure around the *fw3.2* region, but this selection is likely to have predated domestication because reduction is found in both *S. pimpinellifolium* and *S. l. lycopersicum*. Exploration of possible origin of M9 mutation using a wide array of tomato accessions showed that the increase in the derived allele frequency followed the same path as the proposed domestication of tomato and was consistent with the selection for a larger fruit along this process (5). However, the allele did not become fixed early during domestication of tomato, further supporting the notion that this region may not have been under strong selection during domestication. Even though the derived M9 allele was detected in low frequency in the Peruvian *S. pimpinellifolium*, without further study we cannot ascertain the validity of this finding and whether the allele may have originated in a wild relative.

Conclusion

To date, only one gene underlying a fruit mass QTL is known (10). Our study identified *SIKLUH* as the second major gene controlling fruit mass in a vegetable or fruit crop, shedding light on the molecular regulation of this trait in plants. Additionally, previous studies had shown an overlap of the *fw2.2/cnr* QTL in chile pepper, albeit that the effect of the pepper QTL was minor (7, 41), whereas the existence of a pepper *fw3.2* QTL was less certain. The findings from this study raise the possibility that *KLUF* may regulate fruit mass in chile pepper and potentially other crops.

Materials and Methods

Plant Materials and Phenotypic Evaluation. Plants were grown under field and greenhouse condition at Ohio Agricultural Research and Development Center (Wooster, OH) in 2010–2012, and a subset in Gainesville, FL, in 2011. Most experiments were conducted with 10 plants for *fw3.2(ys)* and *fw3.2(wt)* NILs and multiple independent lines for the *ORF6*-RNAi and *ORF7*-amiRNA constructs. Fruit and seed mass analyses in *fw3.2* NILs, *ORF6*-RNAi and *ORF7*-amiRNA lines were performed with 20 ripe fruits per plant, unless fruit set was impaired. Side shoot number and lengths were measured at 56 and 70 d after sowing, whereas plant height and number of nodes were recorded at 35 and 56 d after sowing. Leaf attributes were measured on 12th and 13th leaves. Floral organ measurements were performed with five flowers per plant. For cell measurement of the ovaries at anthesis, three ovaries per plant and three sections per ovary were analyzed. For cell measurements at mature fruit stage, two fruits per plant and two sections per fruit were analyzed. To test source–sink relationship, a total of 12 fruits per plant were kept and fruit weight from these plants was compared with control plants with no fruit removal. All phenotypic evaluations were performed independently at least twice, except the fruit removal experiment, which was done once.

Whole Fruit, Pericarp, and Cell Measurements. Mature fruit perimeter, area of pericarp, septum, columella, and placenta were analyzed using Tomato Analyzer (version 2.3) (42). For the cell measurements in the pericarp, thick sections were made with a razor blade, stained with 0.5% Toluidine Blue in 0.1% Sodium Carbonate solution (SPI, Electron Microscopy Supplies), and photographed with a Leica MZFLIII dissecting microscope coupled to a digital camera (SPOT RT KE, Diagnostic Instruments), followed by analysis with ImageJ software. For cell measurements of the ovary at anthesis, the tissue

was fixed in 0.1 M phosphate buffer (pH 7.2) containing 3% (wt/vol) glutaraldehyde and 2% (wt/vol) paraformaldehyde. Samples were dehydrated in 25–90% ethanol series, followed by infiltration in 50–100% (vol/vol) LR White resin (Electron Microscopy Supplies) gradient at room temperature. Samples were embedded in 100% LR White for 24 h at 55 °C. For each ovary, 5-μm transverse sections were stained and photographed and cell measurements were similar as described for the pericarp cell measurements.

Fine Mapping, Gene Action, and Genetic Segregation Analysis. To identify additional recombinants, 178 seedlings of 09569-68 line were screened, resulting in recombinant line 105187-82. The progeny testing of the other line was from a previous study (14). For the genetic segregation analysis of the M9 SNP, we developed five *F*₂ and BC₁*F*₂ populations each that were derived from parents from the core collection showing few polymorphisms in addition to M9. Alleles for the other known fruit weight/shape genes that were segregating in the populations were fixed such that only *fw3.2* was segregating. For the progeny testing and segregation analysis, 10–13 plants per genotype [genotypes carrying homozygous *fw3.2(ys)* and homozygous *fw3.2(wt)* alleles] were grown and 20 ripe fruits representing the average per plant were weighed. The average fruit mass of each plant was contrasted between the genotypes and the significance of fruit mass segregation in each family was determined using Student *t* test. For gene action analysis, 10 plants each for *fw3.2(ys)*, *fw3.2(wt)*, and heterozygous for *fw3.2* were analyzed. Gene action was represented as *D/A*, where *D* = *Aa* – (*AA* + *aa*)/2 and *A* = (*AA* – *aa*)/2. Gene action experiments were repeated with the same NIL family grown in two separate fields.

Gene Expression Analysis. Total RNA was extracted using TRIzol (Invitrogen) as recommended by the manufacturer. Northern blots were performed as described (43). For analyzing *ORF7* transcript accumulation, total RNA was isolated from mature leaves, genomic DNA was removed using TURBO DNA-free kit (Invitrogen), followed by first-strand cDNA synthesis and quantitative PCR (qPCR) analysis using iQSYBR Green Supermix (Bio-Rad). For Northern blot and qPCR analyses, the expression of *elF4α* and *CAC* were used as internal control (for primers, see Table S7). For RNA seq analysis, strand-specific libraries of ~250-bp fragments were prepared using 10 μg of total RNA. Four replicates were generated for each tissue type and time point. Single end sequences of 51 bp were generated on an Illumina HiSeq2000 at Weill Medical College (44). Datasets for the digital gene expression and transcriptome analysis of developing fruit using RNA seq have been published previously, in addition to those published herein (45). Analyzed data can be accessed at the Tomato Functional Genomics Database (<http://ted.bti.cornell.edu/cgi-bin/TFGD/digital/experiment.cgi?ID=D006> and <http://ted.bti.cornell.edu/cgi-bin/TFGD/digital/experiment.cgi?ID=D008>).

Association of *CaKLUH* with Fruit Weight in Chile Pepper. To identify nucleotide polymorphisms, the 5' upstream, intron, 3' downstream regions of the *CaKLUH* ortholog in pepper were sequenced in *C. annuum* cv. "NuMex RNaky" and the *C. frutescens* accession BG 2814-6 (46). Genomic DNA was extracted from 106 individuals from a *F*₃ RIL population derived from a cross between these parents. The intron SNP was developed into a CAPS marker using the primer pair *fw3.2_pepper_F* and *fw3.2_pepper_R* and restriction enzyme digestion with *Hae*III. Ten fruit were dried and weighed from each line.

Development of Transgenic RNA Interference Lines for *ORF6* and *ORF7*. Two RNAi hairpin constructs for *ORF6*, pMC2 corresponding to RNAi-3 transgenic lines and pMC3 corresponding to RNAi-2 transgenic lines, were generated. RNAi-2 targets the coding sequence of *SIKLUH*, whereas RNAi-3 targets mainly the 3'UTR (Fig. 2B). Sense and antisense arms were cloned in the *Hind*III-*Xho*I and *Sac*I-*Xba*I sites of a modified pKYLX80 vector, where *Xho*I and *Sac*I sites are separated by a 150-bp ω-3 fatty acid desaturase intron. Expression cassettes between *Hind*III-*Xba*I were subcloned into binary vector pKYLX71 under a 35S promoter with duplicated enhancer (47). Four artificial miRNAs specific for *ORF7* were designed using WMD3- Web MicroRNA Designer (48) and were cloned into pKYLX71 binary vector. Primers for the RNAi and sequences for amiRNAs are shown in Table S7. All constructs were introduced in the *fw3.2(ys)* NIL 085591-10 carrying an introgression of ~130 kb. Transformations were carried out at the Plant Transformation Core Research Facility at the University of Nebraska-Lincoln.

Association Mapping. For the association mapping, ~7 kb of the *fw3.2* locus was sequenced and additional markers were used to genotype the region in 86 accessions of the core collection (20). Genomic positions and scores of markers used in association analysis are presented in Table S3. Association analysis was done using MLM model of TASSEL2.1 software (49). Q and

K matrix were generated with STRUCTURE 2.2 (50) and SPAGeDi (51), respectively. To generate Q and K matrix, 20 EST-simple sequence repeat markers distributed throughout the genome were used and the data for these markers in the core collection was obtained from a previous study (20).

Molecular Diversity and Phylogenetic Analysis. For the molecular diversity analysis, fragment A (637 bp) and fragment B (507 bp) were sequenced in 157 and 172 lines, respectively (20). Fragment A and B span genomic positions 58850848–58851481 and 58852273–58852778, respectively (positions are based on the Tomato WGS Chromosomes SL2.40). The molecular diversity analysis π and Tajima's D were computed using DnaSP5.0 (52) with a sliding window length 100 and step size 25. Nucleotide diversity across chromosome 3 was calculated using 10 accessions each of *S. l. lycopersicum* and *S. pimpinellifolium*.

A contig of 2,528 bp comprised of five fragments spanning the 4,382-bp region (corresponding to 58848391–58852772 in Tomato WGS chromosome

SL2.40ch03) around *SIKLUH* was obtained from the 86 accessions of the association mapping collection (Dataset S1) and used for the phylogenetic analysis. All sequences were aligned using ClustalX2.1 with default multiple sequence alignment parameters and DNA weight matrix ClustalW (1.6). The alignment file was imported to MEGA5.05 (53) and converted into mega (.meg) file format. The phylogeny was reconstructed using the Neighbor-Joining statistical method with 1,000 bootstrap replications and Maximum Composite Likelihood model. For computing phylogenetic distance, a cut off value of 50% bootstrap value was used.

ACKNOWLEDGMENTS. We thank Dr. D. Choi (Seoul National University) for providing DNA sequence of chile pepper *CaKLUH* region; and Molecular and Cellular Imaging Center, The Ohio State University, for assistance with microscopy. This work was supported by National Science Foundation Grant IOS-0922661 (to E.v.d.K.).

- Darwin C (1868) *The Variation of Animals and Plants Under Domestication* (John Murray, London).
- Diamond J (1997) *Guns, Germs, and Steel* (WW Norton & Co, New York).
- Smith BD (1995) Seed plant domestication in eastern North America. *Last Hunters-First Farmers* 354:193–213.
- Pickersgill B (2007) Domestication of plants in the Americas: Insights from Mendelian and molecular genetics. *Ann Bot (Lond)* 100(5):925–940.
- Blanca J, et al. (2012) Variation revealed by SNP genotyping and morphology provides insight into the origin of the tomato. *PLoS ONE* 7(10):e48198.
- Klee HJ, Giovannoni JJ (2011) Genetics and control of tomato fruit ripening and quality attributes. *Annu Rev Genet* 45:41–59.
- Paran I, van der Knaap E (2007) Genetic and molecular regulation of fruit and plant domestication traits in tomato and pepper. *J Exp Bot* 58(14):3841–3852.
- Tanksley SD (2004) The genetic, developmental, and molecular bases of fruit size and shape variation in tomato. *Plant Cell* 16(Suppl):S181–S189.
- Grandillo S, Ku HM, Tanksley SD (1999) Identifying loci responsible for natural variation in fruit size and shape in tomato. *Theor Appl Genet* 99(6):978–987.
- Frary A, et al. (2000) *fw2.2*: A quantitative trait locus key to the evolution of tomato fruit size. *Science* 289(5476):85–88.
- Guo M, et al. (2010) *Cell Number Regulator1* affects plant and organ size in maize: Implications for crop yield enhancement and heterosis. *Plant Cell* 22(4):1057–1073.
- Libault M, et al. (2010) A member of the highly conserved *FWL* (tomato *FW2.2*-like) gene family is essential for soybean nodule organogenesis. *Plant J* 62(5):852–864.
- van der Knaap E, Tanksley SD (2003) The making of a bell pepper-shaped tomato fruit: Identification of loci controlling fruit morphology in Yellow Stuffer tomato. *Theor Appl Genet* 107(1):139–147.
- Zhang N, Brewer MT, van der Knaap E (2012) Fine mapping of *fw3.2* controlling fruit weight in tomato. *Theor Appl Genet* 125(2):273–284.
- Blas AL, et al. (2012) Genetic mapping of quantitative trait loci controlling fruit size and shape in papaya. *Mol Breed* 29(2):457–466.
- Costantini L, Battilana J, Lamaj F, Fanizza G, Grando MS (2008) Berry and phenology-related traits in grapevine (*Vitis vinifera* L.): From quantitative trait loci to underlying genes. *BMC Plant Biol* 8:38.
- Doganlar S, Frary A, Daunay MC, Lester RN, Tanksley SD (2002) Conservation of gene function in the solanaceae as revealed by comparative mapping of domestication traits in eggplant. *Genetics* 161(4):1713–1726.
- Eduardo I, et al. (2011) QTL analysis of fruit quality traits in two peach intraspecific populations and importance of maturity date pleiotropic effect. *Tree Genet Genomes* 7(2):323–335.
- Zhang GR, et al. (2010) Fruit size QTL analysis of an F-1 population derived from a cross between a domesticated sweet cherry cultivar and a wild forest sweet cherry. *Tree Genet Genomes* 6(1):25–36.
- Ranc N, Muñoz S, Santoni S, Causse M (2008) A clarified position for *Solanum lycopersicum* var. *cerasiforme* in the evolutionary history of tomatoes (solanaceae). *BMC Plant Biol* 8:130.
- Zhang N (2012) Fine mapping and characterization of *fw3.2*, one of the major QTL controlling fruit size in tomato. PhD dissertation (The Ohio State Univ, Columbus, OH).
- Anastasiou E, et al. (2007) Control of plant organ size by *KLUH/CYP78A5*-dependent intercellular signaling. *Dev Cell* 13(6):843–856.
- Paterson AH, et al. (1995) Convergent domestication of cereal crops by independent mutations at corresponding genetic loci. *Science* 269(5231):1714–1718.
- Adamski NM, Anastasiou E, Eriksson S, O'Neill CM, Lenhard M (2009) Local maternal control of seed size by *KLUH/CYP78A5*-dependent growth signaling. *Proc Natl Acad Sci USA* 106(47):20115–20120.
- Fang W, Wang Z, Cui R, Li Y (2012) Maternal control of seed size by *EOD3/CYP78A6* in *Arabidopsis thaliana*. *Plant J* 70(6):929–939.
- Ito T, Meyerowitz EM (2000) Overexpression of a gene encoding a cytochrome P450, *CYP78A9*, induces large and seedless fruit in *Arabidopsis*. *Plant Cell* 12(9):1541–1550.
- Miyoshi K, et al. (2004) *PLASTOCHRON1*, a timekeeper of leaf initiation in rice, encodes cytochrome P450. *Proc Natl Acad Sci USA* 101(3):875–880.
- Katsumata T, et al. (2011) Involvement of the *CYP78A* subfamily of cytochrome P450 monooxygenases in protonema growth and gametophore formation in the moss *Physcomitrella patens*. *Biosci Biotechnol Biochem* 75(2):331–336.
- Meyer KB, et al. (2011) A functional variant at a prostate cancer predisposition locus at 8q24 is associated with PVT1 expression. *PLoS Genet* 7(7):e1002165.
- Näkki A, et al. (2010) Allelic variants of *IL1R1* gene associate with severe hand osteoarthritis. *BMC Med Genet* 11:50.
- Nischwitz S, et al. (2011) More CLEC16A gene variants associated with multiple sclerosis. *Acta Neurol Scand* 123(6):400–406.
- Uno S, et al. (2010) A genome-wide association study identifies genetic variants in the *CDKN2BAS* locus associated with endometriosis in Japanese. *Nat Genet* 42(8):707–710.
- Sandal NN, Bojsen K, Marcker KA (1987) A small family of nodule specific genes from soybean. *Nucleic Acids Res* 15(4):1507–1519.
- Stougaard J, Jørgensen JE, Christensen T, Kühle A, Marcker KA (1990) Interdependence and nodule specificity of cis-acting regulatory elements in the soybean leghemoglobin *lbc3* and *N23* gene promoters. *Mol Gen Genet* 220(3):353–360.
- Vieweg MF, et al. (2004) The promoter of the *Vicia faba* L. leghemoglobin gene *VfLb29* is specifically activated in the infected cells of root nodules and in the arbuscule-containing cells of mycorrhizal roots from different legume and nonlegume plants. *Mol Plant Microbe Interact* 17(1):62–69.
- Gillaspay G, Ben-David H, Grussem W (1993) Fruits: A developmental perspective. *Plant Cell* 5(10):1439–1451.
- Iglesias DJ, et al. (2007) Physiology of citrus fruiting. *Braz J Plant Physiol* 19(4):333–362.
- Marcelis L, Hofman-Eijer L (1993) Cell division and expansion in the cucumber fruit. *J Hortic Sci* 68(5):665–671.
- Xiao H, et al. (2009) Integration of tomato reproductive developmental landmarks and expression profiles, and the effect of *SUN* on fruit shape. *BMC Plant Biol* 9:49.
- Cong B, Liu J, Tanksley SD (2002) Natural alleles at a tomato fruit size quantitative trait locus differ by heterochronic regulatory mutations. *Proc Natl Acad Sci USA* 99(21):13606–13611.
- Rao GU, Ben Chaim A, Borovsky Y, Paran I (2003) Mapping of yield-related QTLs in pepper in an interspecific cross of *Capsicum annuum* and *C. frutescens*. *Theor Appl Genet* 106(8):1457–1466.
- Brewer MT, et al. (2006) Development of a controlled vocabulary and software application to analyze fruit shape variation in tomato and other plant species. *Plant Physiol* 141(1):15–25.
- Xiao H, Jiang N, Schaffner E, Stockinger EJ, van der Knaap E (2008) A retrotransposon-mediated gene duplication underlies morphological variation of tomato fruit. *Science* 319(5869):1527–1530.
- Zhong S, et al. (2011) High-throughput illumina strand-specific RNA sequencing library preparation. *Cold Spring Harb Protoc* 2011(8):940–949.
- Huang J, Van Houten J, Gonzalez G, Xiao H, van der Knaap E (2013) Genome-wide identification, phylogeny and expression analysis of *SUN*, *OFP* and *YABBY* gene family in tomato. *Mol Genet Genomics* 288(3-4):111–129.
- Ben-Chaim A, et al. (2006) QTL analysis for capsaicinoid content in *Capsicum*. *Theor Appl Genet* 113(8):1481–1490.
- Schardl CL, et al. (1987) Design and construction of a versatile system for the expression of foreign genes in plants. *Gene* 61(1):1–11.
- Ossowski S, Schwab R, Weigel D (2008) Gene silencing in plants using artificial microRNAs and other small RNAs. *Plant J* 53(4):674–690.
- Bradbury PJ, et al. (2007) TASSEL: Software for association mapping of complex traits in diverse samples. *Bioinformatics* 23(19):2633–2635.
- Falush D, Stephens M, Pritchard JK (2007) Inference of population structure using multilocus genotype data: Dominant markers and null alleles. *Mol Ecol Notes* 7(4):574–578.
- Hardy OJ, Vekemans X (2002) SPAGeDi: A versatile computer program to analyse spatial genetic structure at the individual or population levels. *Mol Ecol Notes* 2(4):618–620.
- Librado P, Rozas J (2009) DnaSP v5: A software for comprehensive analysis of DNA polymorphism data. *Bioinformatics* 25(11):1451–1452.
- Tamura K, et al. (2011) MEGA5: Molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol Biol Evol* 28(10):2731–2739.