

REVIEW ARTICLE

Open Access

Perspectives of CRISPR/Cas-mediated *cis*-engineering in horticulture: unlocking the neglected potential for crop improvement

Qiang Li^{1,2}, Manoj Sapkota³ and Esther van der Knaap^{2,3,4}

Abstract

Directed breeding of horticultural crops is essential for increasing yield, nutritional content, and consumer-valued characteristics such as shape and color of the produce. However, limited genetic diversity restricts the amount of crop improvement that can be achieved through conventional breeding approaches. Natural genetic changes in *cis*-regulatory regions of genes play important roles in shaping phenotypic diversity by altering their expression. Utilization of CRISPR/Cas editing in crop species can accelerate crop improvement through the introduction of genetic variation in a targeted manner. The advent of CRISPR/Cas-mediated *cis*-regulatory region engineering (*cis*-engineering) provides a more refined method for modulating gene expression and creating phenotypic diversity to benefit crop improvement. Here, we focus on the current applications of CRISPR/Cas-mediated *cis*-engineering in horticultural crops. We describe strategies and limitations for its use in crop improvement, including de novo *cis*-regulatory element (CRE) discovery, precise genome editing, and transgene-free genome editing. In addition, we discuss the challenges and prospects regarding current technologies and achievements. CRISPR/Cas-mediated *cis*-engineering is a critical tool for generating horticultural crops that are better able to adapt to climate change and providing food for an increasing world population.

Introduction

Horticultural crops comprise vegetables, fruits, and ornamental flowers as well as aromatic and medicinal plants, thereby providing essential resources to society. For example, the availability and consumption of a wide variety of vegetables and fruits allow us to meet our daily dietary needs. Moreover, we enlighten our days with the abundance of floriculture products for aesthetic uses and visual enjoyment. Collectively, horticultural crops make essential contributions to humankind while also providing the economic engines that drive the success of societies all over the world¹.

Despite their collective importance, the improvement of many horticultural crops has lagged behind most agronomic crops, such as rice, corn, and soybean. Yet, improvement of horticultural crops for traits such as resistance to biotic and abiotic stresses, yield, and health-related nutrients would benefit the entire sector. Genetic diversity is a critical source for crop improvement. However, this diversity is often limiting, especially for certain species². The limited genetic diversity could result in significant obstacles for further improvement by conventional breeding approaches. Research in several crops has demonstrated that much of the genetic changes underlying traits of economic importance reside in the *cis*-regulatory regions of genes^{3,4}. These changes appear to have been selected during domestication, resulting in desirable traits caused by altered gene expression^{3,5}. The CRISPR/Cas-based platform offers a powerful tool by engineering *cis*-regulatory regions (*cis*-engineering) to introduce genetic diversity that could potentially

Correspondence: Esther van der Knaap (vanderkn@uga.edu)

¹College of Horticultural Science and Engineering, Shandong Agricultural University, Tai'an, China

²Center for Applied Genetic Technologies, University of Georgia, Athens, GA, USA

Full list of author information is available at the end of the article.

© The Author(s) 2020



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

accelerate crop improvement^{6–10}. Despite the importance of regulatory changes in genes, the application of CRISPR/Cas-mediated *cis*-engineering has only been explored sporadically. The genome sequence for at least 181 horticultural species is available¹¹ and genome editing has been used to generate primarily knockout mutations in at least 25 of them^{12–17}. These achievements demonstrate the feasibility of applying CRISPR/Cas-mediated *cis*-engineering to expand the phenotypic diversity of many horticultural crops.

Natural variation in *cis*-regulatory regions resulting from the domestication of horticultural crops

Cis-regulatory regions are non-coding DNA sequences that control the transcription of genes¹⁸. These *cis*-regulatory sequences consist of combinations of CREs that affect gene expression level often in a spatiotemporal manner^{9,18,19}. Single-nucleotide polymorphisms (SNPs), insertions, deletions, inversions, and epigenetic variations are the most common natural variation in *cis*-regulatory regions that are associated with domestication. Some examples from horticultural crops are discussed below.

Single-nucleotide polymorphisms

Genomic studies in horticultural crops have generated insights into the role of SNP in shaping phenotypic diversity among individuals²⁰. During tomato (*Solanum lycopersicum*) domestication, selection frequently occurred for fruit size and shape, traits that show extensive variation and large increases over that of the wild relatives⁴. Increases in fruit weight are thought to be controlled by SNPs in the promoter of *FW2.2* (*SICNR*) and *FW3.2* (*SIKLUH*)^{4,21,22}. The *lc* allele contains two SNPs in a 15-bp repressor element downstream of tomato *WUSCHEL* (*SIWUS*). The SNPs are proposed to prevent the binding of the MADS-box transcription factor AGAMOUS, which is required to recruit the repressive Polycomb proteins to shut down *SIWUS* expression, thereby ultimately resulting in larger fruits^{4,23,24}. In another example in tomato, two SNPs in the promoter of *Slcyc-B* are highly associated with high β -carotene content²⁵.

In citrus (*Citrus reticulata*), a recent report found an SNP in a miniature inverted-repeat transposable element (MITE) in the promoter of *carotenoid cleavage dioxygenase 4b* (*CCD4*) to be sufficient to increase the expression of this gene, resulting in red coloration of fruit peel²⁶. In pepper (*Capsicum chinense*), an SNP in the promoter of *MYB31* is associated with a hyperfunctional W-box, leading to stronger binding of WRKY9. This stronger binding is associated with enhanced expression of *MYB31*, resulting in extremely pungent peppers²⁷.

Insertions

Insertions are sources of genetic diversity that can alter gene expression by introducing new or disrupting existing CREs. Especially transposable elements (TEs) play important roles in creating genomic variation by altering gene regulation^{28,29}. TE-induced variations in *cis*-regulatory region are also important in the shaping of domestication-related phenotypes in many horticultural crops. One example is the tomato fruit shape gene *SUN*. The transposition event at the *sun* locus mediated by the *Rider* retrotransposon placed a copy of *SUN* in addition to *Rider* itself in the intron of *DEFL1*. The ancestral copy of *SUN* on chromosome 10 is lowly expressed, but its derived copy on chromosome 7, where the *sun* locus maps, is highly expressed³⁰. The high expression of *SUN* on chromosome 7 is thought to be from the promoter of *DEFL1* that would now serve as an enhancer of *SUN*, leading to the elongated tomato fruit³¹. Another *Rider* insertion in the first intron of *SEPALLATA4* (*SEP4*) leads to a jointless pedicel, reduced fruit dropping, which facilitates mechanical harvesting³². In grape (*Vitis vinifera*), the insertion of the *Gret1* (Grapevine Retrotransposon 1) in the *VvMYBA1* promoter leads to its inactivation, resulting in a white berry phenotype³³. In blood oranges (*Citrus sinensis*), the insertion of a *Copia*-like retrotransposon controls the expression of *Ruby* and the cold dependency of anthocyanin production in the fruit³⁴. In cauliflower (*Brassica oleracea* var *botrytis*), a 695-bp *Harbinger* DNA transposon insertion in the *MYB2* promoter increases the expression of this gene, resulting in a purple phenotype³⁵. Additionally, the differentiation of winter and spring genotypes in rapeseed (*Brassica napus* L.) primarily arose from a MITE transposon insertion in the upstream region of *BnFLC.A10*³⁶.

Other examples of insertions that are possibly associated with TE activity are found as well. In tomato, *ej2^w* (*enhancer-of-jointless 2*) is a weak loss-of-function allele, which was selected during tomato domestication and caused by a 564-bp insertion in the fifth intron of *EJ2*. The mutation results in unbranched inflorescences with exceptionally large sepals³². An 8-bp insertion in the promoter of *SibHLH59* significantly increased its expression in accessions producing high ascorbic acid levels³⁷. In apple (*Malus × domestica*), multiple repeats of a 23-bp motif in the promoter of *MYB10* generate an auto-regulatory locus, which is sufficient to account for increased expression and ectopic accumulation of anthocyanins in red-fleshed apples³⁸. Another example from apple is that a 36-bp insertion in *MdSAUR37* promoter contributed to high fruit malate content³⁹. In cucumber (*Cucumis sativus* L.), a 10-bp fragment was replaced by an 812-bp fragment in the promoter of *CsHDZIV11/CsGGL3* at the *few spines 1* (*fs1*) locus, giving

rise to higher expression of *CsGL3* and fewer fruit spines⁴⁰.

Deletions and inversions

Deletions are common genetic changes that provide a wealth of domesticated related phenotypic diversity. One remarkable example is a 31-kb deletion upstream of tomato *OVATE Family Protein 20* (*SIOFP20*). The deletion is associated with reduced expression of *SIOFP20* and contributes to natural fruit shape variation in the tomato germplasm⁴¹. A 3-bp deletion in the promoter of tomato *Al-ACTIVATED MALATE TRANSPORTER9* (*Sl-ALMT9*) was selected during tomato domestication. The deletion disrupts the repression of *Sl-ALMT9* by *Sl-WRKY42*. This results in increased *Sl-ALMT9* gene expression levels, thereby conferring high fruit malate contents and aluminum tolerance in tomato⁴². Flowering time is an important trait for cucumber domestication. A 39.9-kb deletion and a 16.2-kb deletion located 16.5-kb upstream of cucumber *FLOWERING LOCUS T* (*CsFT*) are both associated with higher *CsFT* expression levels and earlier flowering⁴³. The *CsFT* locus was selected during cucumber domestication and has been important in its adaptation to higher latitudes for cultivation⁴³. Therefore, deletions can confer desirable traits through either decreased gene expression by removing enhancers and binding sites of activators or increased gene expression by removing repressors and binding sites of repressors.

Genomic inversions also play a role in plant domestication as they could have widespread *cis*-regulatory effects⁴⁴. One of the remarkable examples of variation in locule number is controlled by a nearly 300-kb inversion of the *fasciated* (*fas*) locus in tomato. The *fas* locus is characterized by disruption of the promoter region of tomato *CLAVATA3* (*SICLV3*), leading to downregulation of the gene and larger fruit with increased number of locules^{24,45}.

Epigenetic variations

Natural epigenetic variations contribute to heritable phenotypic diversity that is not caused by modification in the DNA sequence^{46–49}. One of the best examples of an epiallelic variant that impacts an important agronomical trait is the *Colorless Non-Ripening* (*Cnr*) allele in tomato. The epiallele of *LeSPL-CNR* is responsible for colorless fruits with a substantial loss of cell-to-cell adhesion⁵⁰. In *Cnr* mutants, hyper-methylation was found along a 286-bp CRE located ~2.4-kb upstream from the first ATG of *LeSPL-CNR*. This change in methylation status likely explains the reduced expression level of *LeSPL-CNR* and the ripening defects in *Cnr* fruits⁵⁰. Another epigenetic mutation was found in the promoter of the tomato *SITAB2* gene. The mutation controls pigment production in tomato leaves that are affected by DNA methylation

level in the promoter of the gene⁵¹. *Vitamin E 3* (*VTE3*) is another naturally occurring epiallele controlling vitamin E accumulation in tomato fruits⁵². The *VTE3* expression in fruits is regulated by DNA methylation in the promoter region of the gene⁵². Additional examples include the control of anthocyanin accumulation in apple and pear (*Pyrus communis*) fruit skin^{53–55} and sex determination in melon (*Cucumis melo*)⁵⁶. There is also increasing evidence that promoter DNA methylation plays an important role in regulating tomato fruit ripening^{57,58}. Notably, the tomato *DML2* is critical for tomato fruit ripening by mediating DNA hypomethylation in promoters of hundreds of genes during development⁵⁸.

Taken together, these studies highlight the importance of genetic and epigenetic divergence in *cis*-regulatory regions, including the upstream regions, introns, and downstream regions of genes. Therefore, natural genetic variants, epialleles, and functional CREs in *cis*-regulatory regions are excellent genome editing targets to create novel variants for the improvement of horticultural crops.


Recent progress in CRISPR/Cas-mediated *cis*-engineering in plants

So far, the most frequent application of CRISPR/Cas has been to target coding sequences with the goals to create null alleles^{59–62}. Although this application greatly facilitates heritable alleles for reverse genetics studies, selection of loss-of-function mutations in coding regions may result in pleiotropic or deleterious effects^{45,63,64}. Compared to coding sequences, modulating gene expression by *cis*-engineering is more likely to benefit crop improvement with less detrimental pleiotropic effects^{3,7,9,10,59,64}.

To date, at least 15 articles described successful CRISPR/Cas-mediated *cis*-engineering via genome editing for 17 genes in eight plants species, including eight genes in four horticultural crops (Fig. 1a). In addition, CRISPR/Cas-mediated *cis*-engineering has also been achieved to edit the epigenome. However, only a handful of cases have been described in *Arabidopsis* that show epigenome editing by altering DNA methylation^{65,66} and histone acetylation⁶⁷. Because of the few examples in epigenome editing, the following sections will only describe the applications of *cis*-engineering of DNA.

Promoter disruption

In tomato, a multiplexed CRISPR/Cas9 promoter targeting approach was used to edit the promoters of genes that control fruit size, inflorescence branching, and plant architecture⁷. Importantly, this approach did neither exploit nor require prior knowledge regarding the structure of promoters and other regulatory sequences. Therefore, the multiplexed CRISPR/Cas9 promoter targeting approach is generally applicable for diverse genes



and traits in many crops. Notably, a CRISPR/Cas9-driven sensitized genetic screen approach can recover a collection of *cis*-regulatory alleles with a continuum of phenotypic effects⁷ (Fig. 1b), providing an avenue for expanding genetic diversity in crops.

CRE disruption/deletion

Functional CREs in *cis*-regulatory regions are obvious targets for expanding genetic diversity. However, only a handful of cases have been reported in plants, in which the CRE disruption/deletion was successfully applied to regulate target gene expression.

The rice *RAV2* gene is transcriptionally regulated by high salinity. CRISPR/Cas-mediated *cis*-engineering was used to target the GT-1 element in the promoter of *OsRAV2* and the results strongly indicate that the GT-1 element controls the salt response of this gene⁶⁸. In barley (*Hordeum vulgare*), the promoter of *HvPAPhy_a* was targeted for three CREs, namely GCN4, Skn1, and RY⁶⁹. The lines with mutations in the targeted region show a significant reduction in phytase activity, indicating the

importance of these CREs for the expression of the gene. Similarly, the deletion of a 149-bp regulatory fragment containing a transcription-activator-like effector (TAL)-Binding Element (EBE) in the promoter of *SUGARS WILL EVENTUALLY BE EXPORTED TRANSPORTERS 11* (*SWEET11*) improved rice disease resistance without affecting rice fertility⁶⁴ (Fig. 1c). This result is advantageous compared to the knockout mutant of *OsSWEET11* that showed a sterile phenotype, which is obviously undesirable in crop improvement. Recently, simultaneously editing of EBEs in the promoters of *SWEET* genes resulted in rice lines with broad-spectrum bacterial blight resistance^{70,71}. Three recent studies in Duncan grapefruit (*Citrus paradisi* Macf.) and Wanjincheng orange (*Citrus sinensis* Osbeck) reported that canker-resistant plants were created through CRISPR/Cas editing of the PthA4 effector binding CREs in the promoter of *LATERAL ORGAN BOUNDARIES 1* (*LOB1*)^{72–74}.

The CRISPR/Cas-mediated *cis*-engineering was also utilized to modify known CREs in introns and downstream of genes. The disruption of the CARg element,

including the two causative SNPs downstream of *SIWUS*, is one of the remarkable examples recreating gain-of-function alleles^{7,75} (Fig. 1c). In *Arabidopsis*, a CTCTGYTY motif in the intron of *YUCCA3* (*YUC3*) was identified by chromatin immunoprecipitation-sequencing (CHIP-seq) and is crucial for recruiting RELATIVE OF EARLY FLOWERING 6 (REF6) to its target loci^{76–78}. The deletion of four repeats of this motif leads to diminished binding of REF6 at the mutant loci. In addition, a 450-bp CRE in the second intron of *Arabidopsis* *AGAMOUS* (*AG*) was deleted by CRISPR/Cas9 and verified as the activator of *AG* gene expression. The deletion of this CRE resulted in early flowering because of a 40% decrease in its expression⁷⁹.

Promoter insertion/swapping

Promoter insertion and swapping can be achieved by homology-directed repair (HDR) with potentially great importance to crop improvement (Fig. 1d). However, HDR has been challenging due to its low efficiency in higher plants^{60,80}. So far, only three cases have been reported, in which the promoters were accurately inserted or swapped by CRISPR/Cas9-mediated HDR^{81–83}. A 35S promoter was inserted upstream of *anthocyanin 1* (*ANT1*), resulting in enhanced anthocyanin accumulation and intensely purple tomato tissues⁸¹. In maize, the HDR pathway was used to insert as well as swap the native *GOS2* promoter in the 5'-untranslated region of *ARGOS8* (Fig. 1d). The edited plants showed increased expression of *ARGOS8* and higher grain yield under drought stress conditions in field trials⁸². Additionally, glyphosate-tolerant cassava (*Manihot esculenta*) was generated by a promoter swap of the *5-enolpyruvylshikimate-3-phosphate synthase* (*EPSPS*) gene⁸³.

These encouraging achievements show the potential for using CRISPR/Cas-mediated *cis*-engineering to improve crop yield, quality, and stress resistance.

Strategies for application of CRISPR/Cas-mediated *cis*-engineering in horticultural crops improvement

De novo CRE discovery

Prior knowledge of CREs in *cis*-regulatory region is helpful to apply *cis*-engineering in crop improvement. Many previously described CREs, especially transcription factor-binding sites (TFBSs), in plant promoters can be identified by submitting sequences to various databases, including TRANSFAC⁸⁴, PLACE⁸⁵, PlantCARE⁸⁶, JASPAR Core PLANTAE⁸⁷, PlantTFDB⁸⁸, and Plant Regulomics⁸⁹. After the TFBSs have been predicted, the regions can be validated by either *in vitro* methods based on DNA–protein interaction, such as DNA electrophoretic mobility shift assay, DNA pull-down and yeast one-hybrid assays, or *in vivo* CHIP-based methods, for


example, CHIP with DNA microarray (CHIP-chip) and CHIP-seq.

However, the vast majority of CREs are unknown or poorly characterized, highlighting the pressing need for *de novo* CRE discovery. The availability of genomic and transcriptomic data for many horticultural crops allows the identification of novel CREs using bioinformatics-based and experimental approaches^{11,90,91}. The *de novo* CRE discovery is based on sequence conservation that exists among groups of genes that are co-expressed as well as gene families within a single genome, and among orthologs of multiple species^{91–93}.

Genes that show similar expression patterns or are in the same gene family are likely to be tightly co-regulated and/or functionally related. Therefore, clustering co-expressing genes and identification of gene families are helpful to explore conserved CREs and uncover their functions for transcriptional regulation. The shared CREs can be identified by the well-established methods such as multiple EM for motif elicitation (MEME)^{94,95} and exhaustive evaluation of matrix motifs (XX motif)^{96,97}. An ensemble strategy was used for *de novo* soybean cyst nematode-inducible motif discovery in the upstream regulatory sequences of 18 co-expressed genes⁹⁸. Another strategy to identify conserved CREs is by comparing promoter sequences of orthologous genes from different species. Phylogenetic footprinting and variations of the technique are designed for the CRE discovery approach^{99–103}. mVISTA is a commonly used tool for comparative analysis of genomic sequences¹⁰⁴. The comparison of the *CLV3* promoters in tomato with three other *Solanaceae* species, *S. pennellii*, potato (*S. tuberosum*), and pepper (*C. annuum*) was performed using mVISTA. This resulted in the identification of three putative CREs between tomato and pepper, and four CREs between tomato and potato⁷. Complementary to bioinformatics-based approaches are experimental approaches, for example, deconstructive and reconstructive approaches, by which numerous inducible and tissue-specific CREs are characterized^{90,105}.

Choice of appropriate approach for CRISPR/Cas-mediated *cis*-engineering

CRISPR/Cas-based technologies offer multiple strategies to engineer *cis*-regulatory regions according to the prior knowledge of the target region or given purpose. If no prior knowledge of the target region exists, multiplexed CRISPR/Cas promoter targeting approach can be applied to putative “negative regulators” of the desirable traits by creating a collection of reduced-function alleles (Fig. 1b). In addition, a well-defined promoter can be exchanged with the promoter of the gene of interest to increase expression level or change temporal/spatial expression pattern of the gene (Fig. 1d). For a given CRE in a target region of interest, the CRE can be disrupted or



deleted on the basis of the random indel mutations introduced by non-homologous end joining (NHEJ) repair pathway^{7,64,69,72–75,78,79} (Fig. 1c).

CRISPR/Cas-mediated point mutations and CRE swaps are also important approaches to manipulate gene expression (Fig. 2). Apart from the above-mentioned SNPs that underlied the domestication of crops, numerous studies also documented that single-nucleotide alterations in regulatory sequences can be sufficient to produce substantial effects on gene expression^{106–108}. For example, in soybean, nucleotide mutations in the core and flanking sequences of G-box element lead to both increases and decreases in gene expression in both native and synthetic promoters¹⁰⁹. In apple, the presence of R6

motif, a binding site of *MdMYB10*, in the promoter of *MdMYB10* results in auto-activation of the gene and elevated anthocyanins³⁸. The synthetic promoters of pear *MYB10* and *Arabidopsis MYB75* harboring the R6 motif significantly increase the expression of these genes, leading to elevated anthocyanin levels in transgenic plants of pear and *Arabidopsis*¹¹⁰. Moreover, the insertion of the R6 motif into the promoter of the gene encoding an anthocyanin biosynthetic enzyme flavonoid 3'5'-hydroxylase (F3'5'H) and a vitamin C biosynthesis gene *GDP-L-Galactose Phosphorylase (GGP)* of kiwifruit (*Actinidia eriantha*) altered the anthocyanin profile and increased vitamin C content in a *MYB10*-dependent manner, respectively¹¹⁰. Therefore, the R6 motif can be harnessed to generate new diversity in many horticultural species to increase anthocyanin content (Fig. 2b).

Transgene-free genome editing

Transgene-free genome editing is the preferred choice for the application of *cis*-engineering for crop improvement and commercialization of genome-edited crops. Genome editing with stable expression of CRISPR/Cas DNA involves the integration of the construct into the host genome, raising concerns about undesirable off-target changes and biosecurity^{60,111}. Genetic segregation by selfing or crossing can be used to obtain transgene-free edited plants. Recently, several strategies have been developed to accelerate the removal of transgene components while retaining the desired mutations. These strategies include the integration of fluorescent markers^{112,113}, herbicide-dependent isolation system¹¹⁴, and the programmed self-elimination system¹¹⁵.

An alternative approach for creating transgene-free edited plants is transient expression of CRISPR/Cas DNA as have been reported in many crops, including wheat^{116,117}, barley¹¹⁸, tetraploid potato^{119,120}, tomato¹²¹, and cotton¹²². Compared to stable transformation of CRISPR/Cas DNA, transient expression is especially useful in certain horticultural crops that are vegetatively propagated, self-incompatible, polyploid, and/or have long juvenile stages¹²³.

Given that traditional breeding, including chemically and physically induced mutagenesis, and DNA-based genome editing may introduce off-target mutations, editing in a DNA-free manner via preassembled Cas9 protein-guide RNA (gRNA) ribonucleoproteins (RNPs) is an increasingly popular approach due to higher specificity, and low off-target mutations further alleviating public concerns^{124–127}. RNPs have been adopted in the transformation of protoplasts in some horticultural crops, such as lettuce (*Lactuca sativa* L.)¹²⁸, *petunia*¹²⁹, apple and grape¹³⁰, and potato¹³¹. However, regeneration of mature plants from the edited protoplasts is still challenging for most of the horticultural crops.

Currently available transgene-free genome editing approaches are primarily conducted through traditional transformation methods that require tissue culture, which is a labor-intensive process. Therefore, tissue culture-free methods are highly desirable and necessary for transgene-free genome editing. *In planta* transformation takes advantage of natural biological processes, which makes it a valuable alternative to in vitro tissue culture and regeneration^{132,133}. Various plant cells or tissues can be the ideal transformation targets such as germline or meristematic cells^{116,134,135} and dormant buds¹³⁶. Recently, *in planta* particle bombardment has been used to deliver CRISPR/Cas9 DNA into shoot apical meristems, resulting in transgene-free homozygous mutated wheat plants¹³⁴. Moreover, recent efforts have been made to deliver RNPs into embryo cells in maize¹³⁵ and wheat¹¹⁶ by particle bombardment and into zygotes by polyethylene glycol-Ca²⁺-mediated transfection in rice¹²⁷.

Challenges and prospects

Genome complexity of horticultural crops

The genome sizes of horticultural crops are diverse, ranging from ~200-Mb in some crops, for example, peach (*Prunus persica*), to over 30-Gb in garlic (*Allium sativum*) and onion (*Allium cepa*)¹¹. Many horticultural crops underwent polyploidy, posing extra challenges for genome editing using CRISPR/Cas technologies. Genome editing of polyploid crops requires increased efficiency to edit multiple alleles simultaneously. Even so, CRISPR/Cas technologies have been successfully applied in many polyploid crops due to continuous improvements, including highly active Cas nuclease, multiplex genome editing, and efficient expression systems^{63,137,138}. In case of octoploid and highly heterozygous cultivated strawberry (*Fragaria × ananassa* cv. Camarosa), all five alleles of *FaTM6* were successfully edited using CRISPR/Cas9-mediated dual single-guide (sg) RNA approach¹³⁹. Although the genome of *Fragaria × ananassa* is not yet available, the diploid wild strawberry *F. vesca* reference genome was employed to analyze the allelic variation in the *FaTM6* locus. In this regard, a workflow has been proposed for CRISPR/Cas-mediated mutagenesis for plant species that lack genome sequence information, or feature high heterozygosity or ploidy levels¹⁴⁰. This workflow could be also applicable for many horticultural crops.

High-throughput de novo discovery of CREs in their native context

Currently, experimental validation of predicted CREs largely rely on in vitro techniques that are low accuracy and slow throughput. In recent years, new applications, such as DNase-seq (DNase I hypersensitive sites sequencing), ATAC-seq (assay for transposase-accessible chromatin using sequencing), and CHIP-seq, have significantly


increased our understanding of transcriptional regulatory elements¹⁰⁸. However, these techniques only provide circumstantial evidence and cannot assess the function of CREs in their native context¹⁰⁸. As a complementary approach, CRISPR/Cas-based tiling screen approach was developed in mammalian cells to pinpoint functional CREs in their endogenous context¹⁴¹. The strategy is to densely tile gRNAs across a *cis*-regulatory region to map functional regulatory elements^{142–146}. Although the CRISPR/Cas-based tiling screen approach has not been applied for pinpointing CREs at a large scale in plants, its feasibility was demonstrated in tomato by Rodríguez-Leal et al.⁷.

Efficient and precise genome editing

Efficient precise genome editing is required to achieve *cis*-engineering at the nucleotide level. Base editors, including cytidine base editors (CBEs) and adenine base editors (ABEs), are efficient tools for introducing base substitutions at target sites beyond double-strand breaks^{147,148}. Until now, only CBEs have been optimized and applied for gene function studies in horticultural crops, including tomato^{121,149}, potato^{120,121}, and watermelon¹⁵⁰. Although base editors are good alternatives to HDR-mediated point mutations, it has been challenging to achieve knock-in and replacement of desired CREs in plants. Much efforts has been made to improve the efficiency of HDR through donor design and modulating repair pathways¹³⁸. Recently, a fast and precise multiplexing genome editing method was developed in moss (*Physcomitrella patens*)¹⁵¹. Co-transformation of CRISPR/Cas9 and oligonucleotide templates introduced various mutations into the moss genome with high accuracy and efficiency. It will be interesting to apply such a fast and efficient technology in horticultural crops.

Epigenome editing

The natural epimutations in plants illustrate the potential to further generating phenotypic variation⁴⁶. However, only a small number of natural epialleles have been described in horticultural crops^{50,52–56}. Fortunately, nuclease-dead Cas-mediated epigenome editing technologies hold great promise to expand phenotypic diversity in crops^{46,47}. While some epialleles can be stably inherited over several generations, others epialleles are transient^{50,152–154}. Thus, the stable transmission of editing induced epigenetic changes to the offspring remains unclear^{46,155}. In addition, the expression of CRISPR components may be needed to maintain the trait in the offspring, limiting its application for crop improvement. Further development of CRISPR-based editing tools and the identification of valuable epialleles in horticultural crops will contribute to the application of epigenome editing for expanding phenotypic diversity.



Concluding remarks

We need to continuously improve horticultural commodities to meet the rising demand for food and ornamental production. The widespread applications of CRISPR/Cas technologies in horticultural crops open the possibility for accelerating new variety development^{12–17}. Engineering *cis*-regulatory regions using CRISPR/Cas allows the creation of novel variants, resulting in quantitative variation, and thus holds great potential for creating phenotypic diversity. However, *cis*-engineering is in the beginning stages, and complex relationships between regulation of gene expression by different CREs and the resulting phenotypic changes remains to be fully elucidated⁷. Therefore, using these CRISPR/Cas techniques to screen for desirable traits at the phenotypic level rather than detecting gene expression differences is practical for crop improvement (Fig. 3). Although challenges remain, the application of CRISPR/Cas-mediated *cis*-engineering for horticultural crops improvement will further enhance breeding efforts to improve crop yield, resilience, and commercially desirable traits.

Acknowledgements

Funding in the Van der Knaap laboratory is from the National Science Foundation (IOS 1564366, IOS 1732253, and USDA 2017-67013-26199).

Author details

¹College of Horticultural Science and Engineering, Shandong Agricultural University, Tai'an, China. ²Center for Applied Genetic Technologies, University of Georgia, Athens, GA, USA. ³Institute for Plant Breeding, Genetics and Genomics, University of Georgia, Athens, GA, USA. ⁴Department of Horticulture, University of Georgia, Athens, GA, USA

Author contributions

Q.L. and E.v.d.K. wrote the review. M.S. prepared the figures and edited the manuscript.

Conflict of interest

The authors declare that they have no conflict of interest.

Published online: 15 March 2020

References

- Van den Broeck, G. & Maertens, M. Horticultural exports and food security in developing countries. *Glob. Food Secur.* **10**, 11–20 (2016).
- Shi, J. & Lai, J. Patterns of genomic changes with crop domestication and breeding. *Curr. Opin. Plant Biol.* **24**, 47–53 (2015).
- Swinnen, G., Goossens, A. & Pauwels, L. Lessons from domestication: targeting *cis*-regulatory elements for crop improvement. *Trends Plant Sci.* **21**, 506–515 (2016).
- van der Knaap, E. et al. What lies beyond the eye: the molecular mechanisms regulating tomato fruit weight and shape. *Front. Plant Sci.* **5**, 227 (2014).
- Meyer, R. S. & Purugganan, M. D. Evolution of crop species: genetics of domestication and diversification. *Nat. Rev. Genet.* **14**, 840–852 (2013).
- Li, X., Xie, Y., Zhu, Q. & Liu, Y.-G. Targeted genome editing in genes and *cis*-regulatory regions improves qualitative and quantitative traits in crops. *Mol. Plant* **10**, 1368–1370 (2017).
- Rodríguez-Leal, D., Lemmon, Z. H., Man, J., Bartlett, M. E. & Lippman, Z. B. Engineering quantitative trait variation for crop improvement by genome editing. *Cell* **171**, 470–480 (2017).
- Birchler, J. A. Editing the phenotype: a revolution for quantitative genetics. *Cell* **171**, 269–270 (2017).
- Wolter, F. & Puchta, H. Application of CRISPR/Cas to Understand *Cis*- and *Trans*-Regulatory Elements in Plants. In: *Plant Transcription Factors. Methods in Molecular Biology* (ed. Yamaguchi, N.) vol 1830. (Humana Press, New York, NY, 2018).
- Wolter, F., Schindele, P. & Puchta, H. Plant breeding at the speed of light: the power of CRISPR/Cas to generate directed genetic diversity at multiple sites. *BMC Plant Biol.* **19**, 176 (2019).
- Chen, F. et al. Genome sequences of horticultural plants: past, present, and future. *Hortic. Res.* **6**, 1–23 (2019).
- Xu, J., Hua, K. & Lang, Z. Genome editing for horticultural crop improvement. *Hortic. Res.* **6**, 1–16 (2019).
- Xiong, J.-S., Ding, J. & Li, Y. Genome-editing technologies and their potential application in horticultural crop breeding. *Hortic. Res.* **2**, 15019 (2015).
- Karkute, S. G., Singh, A. K., Gupta, O. P., Singh, P. M. & Singh, B. CRISPR/Cas9 mediated genome engineering for improvement of horticultural crops. *Front. Plant Sci.* **8**, 1635 (2017).
- Koltun, A., Corte, L. E.-D., Mertz-Henning, L. M. & Gonçalves, L. S. Genetic improvement of horticultural crops mediated by CRISPR/Cas: a new horizon of possibilities. *Hortic. Brasil.* **36**, 290–298 (2018).
- Wang, T., Zhang, H. & Zhu, H. CRISPR technology is revolutionizing the improvement of tomato and other fruit crops. *Hortic. Res.* **6**, 1–13 (2019).
- Zhou, J. et al. Application and future perspective of CRISPR/Cas9 genome editing in fruit crops. *J. Integr. Plant Biol.* <https://doi.org/10.1111/jipb.12793> (2019).
- Wittkopp, P. J. & Kalay, G. *Cis*-regulatory elements: molecular mechanisms and evolutionary processes underlying divergence. *Nat. Rev. Genet.* **13**, 59 (2012).
- Bulger, M. & Groudine, M. Functional and mechanistic diversity of distal transcription enhancers. *Cell* **144**, 327–339 (2011).
- Huo, M. A. et al. Identification of functional SNPs in genes and their effects on plant phenotypes. *J. Plant Biotechnol.* **43**, 1–11 (2016).
- Frery, A. et al. fw2.2: a quantitative trait locus key to the evolution of tomato fruit size. *Science* **289**, 85–88 (2000).
- Chakrabarti, M. et al. A cytochrome P450 regulates a domestication trait in cultivated tomato. *Proc. Natl Acad. Sci.* **110**, 17125–17130 (2013).
- Muños, S. et al. Increase in tomato locule number is controlled by two single-nucleotide polymorphisms located near WUSCHEL. *Plant Physiol.* **156**, 2244–2254 (2011).
- Chu, Y. H., Jang, J. C., Huang, Z. & van der Knaap, E. Tomato locule number and fruit size controlled by natural alleles of *lc* and *fas*. *Plant Direct* **3**, e00142 (2019).
- Orchard, C. *Naturally Occurring Variation in the Promoter of the Chromoplast-Specific Cyc-B Gene in Tomato can be used to Modulate Levels of β -Carotene in Ripe Tomato Fruit* (The Ohio State University, 2014).
- Zheng, X. et al. Natural variation in CCD4 promoter underpins species-specific evolution of red coloration in citrus peel. *Mol. Plant* **12**, 1294–1307 (2019).

27. Zhu, Z. et al. Natural variations in the MYB transcription factor MYB31 determine the evolution of extremely pungent peppers. *N. Phytol.* **223**, 922–938 (2019).
28. Lisch, D. How important are transposons for plant evolution? *Nat. Rev. Genet.* **14**, 49 (2013).
29. Olsen, K. M. & Wendel, J. F. A bountiful harvest: genomic insights into crop domestication phenotypes. *Annu. Rev. Plant Biol.* **64**, 47–70 (2013).
30. Xiao, H., Jiang, N., Schaffner, E., Stockinger, E. J. & Van Der Knaap, E. A retrotransposon-mediated gene duplication underlies morphological variation of tomato fruit. *Science* **319**, 1527–1530 (2008).
31. Jiang, N., Gao, D., Xiao, H. & Van Der Knaap, E. Genome organization of the tomato sun locus and characterization of the unusual retrotransposon Rider. *Plant J.* **60**, 181–193 (2009).
32. Soyk, S. et al. Bypassing negative epistasis on yield in tomato imposed by a domestication gene. *Cell* **169**, 1142–1155 (2017).
33. Kobayashi, S., Goto-Yamamoto, N. & Hirochika, H. Retrotransposon-induced mutations in grape skin color. *Science* **304**, 982–982 (2004).
34. Butelli, E. et al. Retrotransposons control fruit-specific, cold-dependent accumulation of anthocyanins in blood oranges. *Plant Cell* **24**, 1242–1255 (2012).
35. Chiu, L.-W. et al. The purple cauliflower arises from activation of a MYB transcription factor. *Plant Physiol.* **154**, 1470–1480 (2010).
36. Hou, J. et al. A Tourist-like MITE insertion in the upstream region of the BnFLC. A10 gene is associated with vernalization requirement in rapeseed (*Brassica napus* L.). *BMC Plant Biol.* **12**, 238 (2012).
37. Ye, J. et al. Genome-wide association analysis identifies a natural variation in basic helix-loop-helix transcription factor regulating ascorbate biosynthesis via D-mannose/L-galactose pathway in tomato. *PLoS Genet.* **15**, e1008149 (2019).
38. Espley, R. V. et al. Multiple repeats of a promoter segment causes transcription factor autoregulation in red apples. *Plant Cell* **21**, 168–183 (2009).
39. Jia, D. et al. Apple fruit acidity is genetically diversified by natural variations in three hierarchical epistatic genes: MdSAUR37, MdPP2CH and MdALMT11. *Plant J.* **95**, 427–443 (2018).
40. Zhang, H. et al. A fragment substitution in the promoter of CsH2DZ11/CsGL3 is responsible for fruit spine density in cucumber (*Cucumis sativus* L.). *Theor. Appl. Genet.* **129**, 1289–1301 (2016).
41. Wu, S. et al. A common genetic mechanism underlies morphological diversity in fruits and other plant organs. *Nat. Commun.* **9**, 4734 (2018).
42. Ye, J. et al. An InDel in the promoter of Al-ACTIVATED MALATE TRANSPORTER9 selected during tomato domestication determines fruit malate contents and aluminum tolerance. *Plant Cell* **29**, 2249–2268 (2017).
43. Wang, S. et al. FLOWERING LOCUS T improved cucumber adaptation to higher latitudes. *Plant Physiol.* **182**, 908–918 (2020).
44. Said, I. et al. Linked genetic variation and not genome structure causes widespread differential expression associated with chromosomal inversions. *Proc. Natl Acad. Sci. USA* **115**, 5492–5497 (2018).
45. Xu, C. et al. A cascade of arabinosyltransferases controls shoot meristem size in tomato. *Nat. Genet.* **47**, 784 (2015).
46. Gallusci, P. et al. Epigenetics for plant improvement: current knowledge and modeling avenues. *Trends Plant Sci.* **22**, 610–623 (2017).
47. Arencibia, A. D., D’Afonseca, V., Chakravarthi, M. & Castiglione, S. Learning from transgenics: advanced gene editing technologies should also bridge the gap with traditional genetic selection. *Electron. J. Biotechnol.* **41**, 22–29 (2019).
48. Kumar, S. Epigenomics for crop improvement: current status and future perspectives. *J. Genet. Cell Biol.* **3**, 128–134 (2019).
49. Sun, S., Wang, X., Wang, K. & Cui, X. Dissection of complex traits of tomato in the post-genome era. *Theor. Appl. Genet.* <https://doi.org/10.1007/s00122-019-03478-y> (2019).
50. Manning, K. et al. A naturally occurring epigenetic mutation in a gene encoding an SBP-box transcription factor inhibits tomato fruit ripening. *Nat. Genet.* **38**, 948 (2006).
51. Gouil, Q., Novák, O. & Baulcombe, D. C. SLTAB2 is the paramutated SUL-FUREA locus in tomato. *J. Exp. Bot.* **67**, 2655–2664 (2016).
52. Quadrana, L. et al. Natural occurring epialleles determine vitamin E accumulation in tomato fruits. *Nat. Commun.* **5**, 4027 (2014).
53. El-Sharkawy, I., Liang, D. & Xu, K. Transcriptome analysis of an apple (*Malus × domestica*) yellow fruit somatic mutation identifies a gene network module highly associated with anthocyanin and epigenetic regulation. *J. Exp. Bot.* **66**, 7359–7376 (2015).
54. Telias, A. et al. Apple skin patterning is associated with differential expression of MYB10. *BMC Plant Biol.* **11**, 93 (2011).
55. Wang, Z. et al. The methylation of the PcMYB10 promoter is associated with green-skinned sport in Max Red Bartlett pear. *Plant Physiol.* **162**, 885–896 (2013).
56. Martin, A. et al. A transposon-induced epigenetic change leads to sex determination in melon. *Nature* **461**, 1135 (2009).
57. Zhong, S. et al. Single-base resolution methylomes of tomato fruit development reveal epigenome modifications associated with ripening. *Nat. Biotechnol.* **31**, 154 (2013).
58. Lang, Z. et al. Critical roles of DNA demethylation in the activation of ripening-induced genes and inhibition of ripening-repressed genes in tomato fruit. *Proc. Natl Acad. Sci. USA* **114**, E4511–E4519 (2017).
59. Pandiarajan, R. & Grover, A. In vivo promoter engineering in plants: are we ready? *Plant Sci.* **277**, 132–138 (2018).
60. Chen, K., Wang, Y., Zhang, R., Zhang, H. & Gao, C. CRISPR/Cas genome editing and precision plant breeding in agriculture. *Annu. Rev. Plant Biol.* **70**, 667–697 (2019).
61. Korotkova, A., Gerasimova, S., Shumny, V. & Khlestkina, E. Crop genes modified using the CRISPR/Cas system. *Russian J. Genet.* **7**, 822–832 (2017).
62. Korotkova, A., Gerasimova, S. & Khlestkina, E. Current achievements in modifying crop genes using CRISPR/Cas system. *Vavilovskii Zh. Genet. Sel.* **8**, 14422 (2019).
63. Morineau, C. et al. Selective gene dosage by CRISPR-Cas9 genome editing in hexaploid *Camelina sativa*. *Plant Biotechnol. J.* **15**, 729–739 (2017).
64. Li, C. et al. A new rice breeding method: CRISPR/Cas9 system editing of the Xa13 promoter to cultivate transgene-free bacterial blight-resistant rice. *Plant Biotechnol. J.* **18**, 313–315 (2019).
65. Gallego-Bartolomé, J. et al. Targeted DNA demethylation of the *Arabidopsis* genome using the human TET1 catalytic domain. *Proc. Natl Acad. Sci.* **115**, E2125–E2134 (2018).
66. Papikian, A., Liu, W., Gallego-Bartolomé, J. & Jacobsen, S. E. Site-specific manipulation of *Arabidopsis* loci using CRISPR-Cas9 SunTag systems. *Nat. Commun.* **10**, 729 (2019).
67. Paixão, J. F. R. et al. Improved drought stress tolerance in *Arabidopsis* by CRISPR/dCas9 fusion with a histone acetyltransferase. *Sci. Rep.* **9**, 8080 (2019).
68. Duan, Y.-B. et al. Identification of a regulatory element responsible for salt induction of rice OsRAV2 through ex situ and in situ promoter analysis. *Plant Mol. Biol.* **90**, 49–62 (2016).
69. Holme, I. B. et al. Evaluation of the mature grain phytase candidate HvPA-Phy_a gene in barley (*Hordeum vulgare* L.) using CRISPR/Cas9 and TALENs. *Plant Mol. Biol.* **95**, 111–121 (2017).
70. Oliva, R. et al. Broad-spectrum resistance to bacterial blight in rice using genome editing. *Nat. Biotechnol.* **37**, 1344–1350 (2019).
71. Xu, Z. et al. Engineering broad-spectrum bacterial blight resistance by simultaneously disrupting variable TALE-binding elements of multiple susceptibility genes in rice. *Mol. Plant* **12**, 1434–1446 (2019).
72. Jia, H., Orbovic, V., Jones, J. B. & Wang, N. Modification of the PthA4 effector binding elements in Type I Cs LOB 1 promoter using Cas9/sg RNA to produce transgenic Duncan grapefruit alleviating XccΔpthA4: dCs LOB 1.3 infection. *Plant Biotechnol. J.* **14**, 1291–1301 (2016).
73. Peng, A. et al. Engineering canker-resistant plants through CRISPR/Cas9-targeted editing of the susceptibility gene Cs LOB 1 promoter in citrus. *Plant Biotechnol. J.* **15**, 1509–1519 (2017).
74. Jia, H., Orbovic, V. & Wang, N. CRISPR-LbCas12a-mediated modification of citrus. *Plant Biotechnol. J.* **17**, 1928–1937 (2019).
75. Li, T. et al. Domestication of wild tomato is accelerated by genome editing. *Nat. Biotechnol.* **36**, 1160–1163 (2018).
76. Cui, X. et al. REF6 recognizes a specific DNA sequence to demethylate H3K27me3 and regulate organ boundary formation in *Arabidopsis*. *Nat. Genet.* **48**, 694 (2016).
77. Li, C. et al. Concerted genomic targeting of H3K27 demethylase REF6 and chromatin-remodeling ATPase BRM in *Arabidopsis*. *Nat. Genet.* **48**, 687 (2016).
78. Li, C. et al. Verification of DNA motifs in *Arabidopsis* using CRISPR/Cas9-mediated mutagenesis. *Plant Biotechnol. J.* **16**, 1446–1451 (2018).
79. Yan, W., Chen, D. & Kaufmann, K. Efficient multiplex mutagenesis by RNA-guided Cas9 and its use in the characterization of regulatory elements in the AGAMOUS gene. *Plant Methods* **12**, 23 (2016).
80. Li, J. et al. Gene replacements and insertions in rice by intron targeting using CRISPR-Cas9. *Nat. Plants* **2**, 16139 (2016).

81. Čermák, T., Baltes, N. J., Čegan, R., Zhang, Y. & Voytas, D. F. High-frequency, precise modification of the tomato genome. *Genome Biol.* **16**, 232 (2015).
82. Shi, J. et al. ARGOS 8 variants generated by CRISPR-Cas9 improve maize grain yield under field drought stress conditions. *Plant Biotechnol. J.* **15**, 207–216 (2017).
83. Hummel, A. W. et al. Allele exchange at the EPSPS locus confers glyphosate tolerance in cassava. *Plant Biotechnol. J.* **16**, 1275–1282 (2018).
84. Matys, V. et al. TRANSFAC[®]: transcriptional regulation, from patterns to profiles. *Nucleic Acids Res.* **31**, 374–378 (2003).
85. Higo, K., Ugawa, Y., Iwamoto, M. & Korenaga, T. Plant *cis*-acting regulatory DNA elements (PLACE) database: 1999. *Nucleic Acids Res.* **27**, 297–300 (1999).
86. Lescot, M. et al. PlantCARE, a database of plant *cis*-acting regulatory elements and a portal to tools for in silico analysis of promoter sequences. *Nucleic Acids Res.* **30**, 325–327 (2002).
87. Mathelier, A. et al. JASPAR 2014: an extensively expanded and updated open-access database of transcription factor binding profiles. *Nucleic Acids Res.* **42**, D142–D147 (2013).
88. Jin, J. et al. PlantTFDB 4.0: toward a central hub for transcription factors and regulatory interactions in plants. *Nucleic Acids Res.* **45**, gkw982 (2016).
89. Ran, X. et al. Plant Regulomics: a data-driven interface for retrieving upstream regulators from plant multi-omics data. *Plant J.* **101**, 237–248 (2019).
90. Mehrotra, R. et al. Designer promoter: an artwork of *cis* engineering. *Plant Mol. Biol.* **75**, 527–536 (2011).
91. Xie, D., Cai, J., Chia, N.-Y., Ng, H. H. & Zhong, S. Cross-species de novo identification of *cis*-regulatory modules with GibbsModule: application to gene regulation in embryonic stem cells. *Genome Res.* **18**, 1325–1335 (2008).
92. Gruel, J., LeBorgne, M., LeMeur, N. & Théret, N. Simple shared motifs (SSM) in conserved region of promoters: a new approach to identify co-regulation patterns. *BMC Bioinform.* **12**, 365 (2011).
93. Burgess, D. & Freeling, M. The most deeply conserved noncoding sequences in plants serve similar functions to those in vertebrates despite large differences in evolutionary rates. *Plant Cell* **26**, 946–961 (2014).
94. Bailey, T. L. & Elkan, C. Fitting a mixture model by expectation maximization to discover motifs in bipolymers. *Proc. Int. Conf. Intell. Syst. Mol. Biol.* **2**, 28–36 (1994).
95. Bailey, T. L. et al. MEME SUITE: tools for motif discovery and searching. *Nucleic Acids Res.* **37**, W202–W208 (2009).
96. Hartmann, H., Guthöhrlein, E. W., Siebert, M., Luehr, S. & Söding, J. *P*-value-based regulatory motif discovery using positional weight matrices. *Genome Res.* **23**, 181–194 (2013).
97. Hickman, R. et al. Architecture and dynamics of the jasmonic acid gene regulatory network. *Plant Cell* **29**, 2086–2105 (2017).
98. Liu, W. et al. Computational discovery of soybean promoter *cis*-regulatory elements for the construction of soybean cyst nematode-inducible synthetic promoters. *Plant Biotechnol. J.* **12**, 1015–1026 (2014).
99. Tagle, D. A. et al. Embryonic ϵ and γ globin genes of a prosimian primate (*Galago crassicaudatus*): nucleotide and amino acid sequences, developmental regulation and phylogenetic footprints. *J. Mol. Biol.* **203**, 439–455 (1988).
100. Thomas, J. et al. Comparative analyses of multi-species sequences from targeted genomic regions. *Nature* **424**, 788 (2003).
101. Liu, B. et al. An integrative and applicable phylogenetic footprinting framework for *cis*-regulatory motifs identification in prokaryotic genomes. *BMC Genomics* **17**, 578 (2016).
102. Van de Velde, J., Heyndrickx, K. S. & Vandepoele, K. Inference of transcriptional networks in *Arabidopsis* through conserved noncoding sequence analysis. *Plant Cell* **26**, 2729–2745 (2014).
103. Yang, J., Chen, X., McDermid, A. & Ma, Q. DMINDA 2.0: integrated and systematic views of regulatory DNA motif identification and analyses. *Bioinformatics* **33**, 2586–2588 (2017).
104. Frazer, K. A., Pachter, L., Poliakov, A., Rubin, E. M. & Dubchak, I. VISTA: computational tools for comparative genomics. *Nucleic Acids Res.* **32**, W273–W279 (2004).
105. Liu, W., Yuan, J. S. & Stewart, C. N. Jr. Advanced genetic tools for plant biotechnology. *Nat. Rev. Genet.* **14**, 781–793 (2013).
106. Canver, M. C. et al. Characterization of genomic deletion efficiency mediated by clustered regularly interspaced palindromic repeats (CRISPR)/Cas9 nuclease system in mammalian cells. *J. Biol. Chem.* **289**, 21312–21324 (2014).
107. Mansour, M. R. et al. An oncogenic super-enhancer formed through somatic mutation of a noncoding intergenic element. *Science* **346**, 1373–1377 (2014).
108. Lopes, R., Korkmaz, G. & Agami, R. Applying CRISPR–Cas9 tools to identify and characterize transcriptional enhancers. *Nat. Rev. Mol. Cell. Biol.* **17**, 597–604 (2016).
109. Zhang, N., McHale, L. K. & Finer, J. J. Changes to the core and flanking sequences of G-box elements lead to increases and decreases in gene expression in both native and synthetic soybean promoters. *Plant Biotechnol. J.* **17**, 724–735 (2019).
110. Brendolise, C. et al. Multiple copies of a simple MYB-binding site confers *trans*-regulation by specific flavonoid-related R2R3 MYBs in diverse species. *Front. Plant Sci.* **8**, 1864 (2017).
111. Hahn, F. & Nekrasov, V. CRISPR/Cas precision: do we need to worry about off-targeting in plants? *Plant Cell Rep.* **38**, 437–441 (2019).
112. Gao, X., Chen, J., Dai, X., Zhang, D. & Zhao, Y. An effective strategy for reliably isolating heritable and Cas9-free *Arabidopsis* mutants generated by CRISPR/Cas9-mediated genome editing. *Plant Physiol.* **171**, 1794–1800 (2016).
113. Aliaga-Franco, N. et al. Identification of transgene-free CRISPR-edited plants of rice, tomato, and arabidopsis by monitoring DsRED fluorescence in dry seeds. *Front. Plant Sci.* **10**, 1150 (2019).
114. Lu, H. P. et al. CRISPR-S: an active interference element for a rapid and inexpensive selection of genome-edited, transgene-free rice plants. *Plant Biotechnol. J.* **15**, 1371 (2017).
115. He, Y. et al. Programmed self-elimination of the CRISPR/Cas9 construct greatly accelerates the isolation of edited and transgene-free rice plants. *Mol. Plant* **11**, 1210–1213 (2018).
116. Zhang, Y. et al. Efficient and transgene-free genome editing in wheat through transient expression of CRISPR/Cas9 DNA or RNA. *Nat. Commun.* **7**, 12617 (2016).
117. Demirer, G. S., Zhang, H., Goh, N. S., González-Grandío, E. & Landry, M. P. Carbon nanotube-mediated DNA delivery without transgene integration in intact plants. *Nat. Protoc.* **14**, 2954–2971 (2019).
118. Gerasimova, S. et al. Targeted genome modification in protoplasts of a highly regenerable Siberian barley cultivar using RNA-guided Cas9 endonuclease. *Vavilovskii Zh. Genet. Sel.* **22**, 1033–1039 (2018).
119. Andersson, M. et al. Efficient targeted multiallelic mutagenesis in tetraploid potato (*Solanum tuberosum*) by transient CRISPR-Cas9 expression in protoplasts. *Plant Cell Rep.* **36**, 117–128 (2017).
120. Veillet, F. et al. The *Solanum tuberosum* GBSSI gene: a target for assessing gene and base editing in tetraploid potato. *Plant Cell Rep.* **38**, 1–16 (2019).
121. Veillet, F. et al. Transgene-free genome editing in tomato and potato plants using agrobacterium-mediated delivery of a CRISPR/Cas9 cytidine base editor. *Int. J. Mol. Sci.* **20**, 402 (2019).
122. Demirer, G. S. et al. High aspect ratio nanomaterials enable delivery of functional genetic material without DNA integration in mature plants. *Nat. Nanotechnol.* **14**, 456 (2019).
123. Chen, L. et al. A method for the production and expedient screening of CRISPR/Cas9-mediated non-transgenic mutant plants. *Horticult. Res.* **5**, 13 (2018).
124. Kouranova, E. et al. CRISPRs for optimal targeting: delivery of CRISPR components as DNA, RNA, and protein into cultured cells and single-cell embryos. *Hum. Gene Ther.* **27**, 464–475 (2016).
125. Liang, Z. et al. Genome editing of bread wheat using biolistic delivery of CRISPR/Cas9 in vitro transcripts or ribonucleoproteins. *Nat. Protoc.* **13**, 413 (2018).
126. Metje-Sprink, J., Menz, J., Modrzejewski, D. & Sprink, T. DNA-free genome editing: past, present and future. *Front. Plant Sci.* **9**, 1065–1080 (2018).
127. Toda, E. et al. An efficient DNA- and selectable-marker-free genome-editing system using zygotes in rice. *Nat. Plants* **5**, 363 (2019).
128. Woo, J. W. et al. DNA-free genome editing in plants with preassembled CRISPR-Cas9 ribonucleoproteins. *Nat. Biotechnol.* **33**, 1162 (2015).
129. Subburaj, S. et al. Site-directed mutagenesis in *Petunia × hybrida* protoplast system using direct delivery of purified recombinant Cas9 ribonucleoproteins. *Plant Cell Rep.* **35**, 1535–1544 (2016).
130. Malnoy, M. et al. DNA-free genetically edited grapevine and apple protoplast using CRISPR/Cas9 ribonucleoproteins. *Front. Plant Sci.* **7**, 1904 (2016).
131. Andersson, M. et al. Genome editing in potato via CRISPR-Cas9 ribonucleoprotein delivery. *Physiol. Plant.* **164**, 378–384 (2018).
132. Niazian, M., Noori, S. S., Galuszka, P. & Mortazavian, S. M. M. Tissue culture-based *Agrobacterium*-mediated and in planta transformation methods. *Soil Water Res.* **53**, 133–143 (2017).
133. Nagle, M. F., Déjardin, A., Pilate, G. & Strauss, S. H. Opportunities for innovation in genetic transformation of forest trees. *Front. Plant Sci.* **9**, 1443 (2018).

134. Hamada, H. et al. Biolistic-delivery-based transient CRISPR/Cas9 expression enables in planta genome editing in wheat. *Sci. Rep.* **8**, 14422 (2018).
135. Svitashv, S., Schwartz, C., Lenderts, B., Young, J. K. & Cigan, A. M. Genome editing in maize directed by CRISPR–Cas9 ribonucleoprotein complexes. *Nat. Commun.* **7**, 13274 (2016).
136. Fujita, K., Matsuoka, T., Suzuki, S. & Takayanagi, T. In planta transformation technique for grapevines (*Vitis vinifera* L.) using dormant buds. *J. Plant Biotechnol. Biotechnol.* **18**, 161–167 (2009).
137. Zaman, Q. U., Li, C., Cheng, H. & Hu, Q. Genome editing opens a new era of genetic improvement in polyploid crops. *Crop J.* **7**, 141–150 (2019).
138. Zhang, Y., Malzahn, A. A., Sretenovic, S. & Qi, Y. The emerging and uncultivated potential of CRISPR technology in plant science. *Nat. Plants* **5**, 778–794 (2019).
139. Martín-Pizarro, C., Triviño, J. C. & Posé, D. Functional analysis of the TM6 MADS-box gene in the octoploid strawberry by CRISPR/Cas9-directed mutagenesis. *J. Exp. Bot.* **70**, 885–895 (2018).
140. Botella, J. R. Now for the hard ones: is there a limit on CRISPR genome editing in crops? *J. Exp. Bot.* **70**, 734–737 (2019).
141. Montalbano, A., Canver, M. C. & Sanjana, N. E. High-throughput approaches to pinpoint function within the noncoding genome. *Mol. Cell* **68**, 44–59 (2017).
142. Canver, M. C. et al. BCL11A enhancer dissection by Cas9-mediated in situ saturating mutagenesis. *Nature* **527**, 192 (2015).
143. Korkmaz, G. et al. Functional genetic screens for enhancer elements in the human genome using CRISPR-Cas9. *Nat. Biotechnol.* **34**, 192 (2016).
144. Rajagopal, N. et al. High-throughput mapping of regulatory DNA. *Nat. Biotechnol.* **34**, 167 (2016).
145. Sanjana, N. E. et al. High-resolution interrogation of functional elements in the noncoding genome. *Science* **353**, 1545–1549 (2016).
146. Diao, Y. et al. A tiling-deletion-based genetic screen for cis-regulatory element identification in mammalian cells. *Nat. Methods* **14**, 629 (2017).
147. Gaudelli, N. M. et al. Programmable base editing of A•T to G•C in genomic DNA without DNA cleavage. *Nature* **551**, 464 (2017).
148. Komor, A. C., Kim, Y. B., Packer, M. S., Zuris, J. A. & Liu, D. R. Programmable editing of a target base in genomic DNA without double-stranded DNA cleavage. *Nature* **533**, 420 (2016).
149. Shimatani, Z. et al. Targeted base editing in rice and tomato using a CRISPR-Cas9 cytidine deaminase fusion. *Nat. Biotechnol.* **35**, 441 (2017).
150. Tian, S. et al. Engineering herbicide-resistant watermelon variety through CRISPR/Cas9-mediated base-editing. *Plant Cell Rep.* **37**, 1353–1356 (2018).
151. Yi, P. & Goshima, G. Fast, efficient, and precise gene editing in the moss *Physcomitrella patens*. *bioRxiv* <https://doi.org/10.1101/643692> (2019).
152. Johannes, F. et al. Assessing the impact of transgenerational epigenetic variation on complex traits. *PLoS Genet.* **5**, e1000530 (2009).
153. Cortijo, S. et al. Mapping the epigenetic basis of complex traits. *Science* **343**, 1145–1148 (2014).
154. Giovannoni, J. Harnessing epigenome modifications for better crops. *J. Exp. Bot.* **67**, 2535–2537 (2016).
155. Hua, K. et al. Perspectives on the application of genome editing technologies in crop breeding. *Mol. Plant* **12**, 1047–1059 (2019).